

Physiology-based versus allometric scaling of clearance in children; an eliminating process based comparison

Andrea N Edginton, Stefan Willmann

Competence Center Systems Biology, Bayer Technology Services GmbH, Leverkusen, Germany

Corresponding author

Andrea Edginton, Systems Biology, Bldg E41, Bayer Technology Services GmbH, 51368 Leverkusen, Germany.
Email: andreanicole.edginton@bayertechnology.com

Aim: To scale clearance from adults to children, allometric and physiology-based approaches have been suggested. The allometric approach is common but does not account for the maturation processes of the eliminating organ and therefore may not be appropriate to scale clearance to young children. The physiology-based approach accounts for maturity but requires detailed knowledge regarding eliminating processes. This study compared the approaches and determined if their appropriate use was dependent on the clearance pathway.

Methods: A data set of experimentally-obtained clearance values was used. Predicted clearances using both approaches were calculated using the mean age and weight of the studied group. The ratio of predicted to observed clearances were graphed separately for compounds with a predominant clearance pathway (> 85%, $n = 8$). Other compounds with multiple clearance pathways were graphed together ($n = 7$). The process specific age at which both

approaches produced accurate clearance predictions was noted.

Results: The allometric approach systematically overestimated clearance in children below a certain age, contrary to the physiology-based approach that accurately predicted clearance at all ages. The age at which clearances were not biased using the allometric equation was within the first year for compounds eliminated via CYP3A4, four years for CYP1A2 and six months for UGT2B7. Allometric scaling systematically overestimated renal clearance in children up to two years and underestimated clearance above two years.

Conclusions: Physiology-based clearance scaling accurately predicted clearance in children from birth to 18 years. The allometric approach is only accurate above the age at which the major clearance pathway is fully, or near to fully, matured. A combined use of approaches is suggested.

Paed Perinat Drug Ther 2006; 7: 146–153

Keywords: clearance scaling – paediatrics – allometric equation – physiology-based

Introduction

Pharmacokinetic studies in children have lagged behind those in adults such that the majority of drugs prescribed to children have not been significantly studied within the paediatric age range¹.

Since the pharmacokinetic profile is an important predictor of the probability of therapeutic or adverse effect events, the profile differences between children and adults are highly important. One of the major drivers of profile differences is the pharmacokinetic parameter of clearance.

Clearance describes the volume of reference fluid (usually plasma) that is removed of compound per unit time. Because of the importance of this parameter to determining the pharmacokinetic profile, *a priori* knowledge of its compound-specific age-related changes would advance the preparation of the paediatric clinical trial.

Predicting clearance in children based on the clearance in an adult has been the topic of recent publications²⁻⁶. A physiology-based approach has been found to be useful for predicting clearance across all age ranges^{2,5}. This has been demonstrated for a number of the major enzymatic (e.g. CYP1A2) and physiological (e.g. glomerular filtration) eliminating processes. The drawback of the approach is that detailed knowledge about the eliminating processes in adults is required and further, the age-dependence of the ontogeny of these clearance processes is needed. Another established approach, allometric scaling, has been demonstrated to determine the age at which the maturity of the processes responsible for clearance no longer affects the clearance value^{7,8}. Thus, the allometric equation can be predictive of clearance in children following a specific age. The approach is only based on the difference in body weight between the child and the adult and has the benefit of being very easy to calculate. Since the approach does not account for maturation processes, it becomes invalid when used below a certain age where the activities of the eliminating processes are not yet fully developed⁵.

The objective of this study was to retrospectively compare the accuracy of the allometric clearance scaling approach with the physiology-based approach using compounds eliminated via various enzymatic and physiological processes (glomerular filtration). Furthermore, by using compounds for which the clearance process in adults is dominated (> 85%) by one clearance pathway, the pathway-specific age range of appropriate use for each scaling approach will be determined.

Methods

Age dependent clearance data set

A data set of observed clearances for children aged premature to 18 years from previous studies was used^{2,9}. Age-dependent observed clearances from 15 compounds were presented. In one previous study², eight of the compounds, where clearance was primarily (> 85%) due to one process, were used to develop ontogeny models for the scaling of clearance via the physiology-based scaling approach. These were gentamicin and isepamicin (renal clearance due to glomerular

filtration), alfentanil and midazolam (CYP3A), caffeine and ropivacaine (CYP1A2) and morphine and lorazepam (UGT2B7). The seven remaining compounds, namely fentanyl, theophylline, paracetamol, ciprofloxacin, lidocaine, levofloxacin and buprenorphine were cleared via multiple pathways (see Table III in Edginton et al² and levofloxacin data in Edginton et al⁹) and were used to evaluate the ontogeny models².

The observed adult clearance value was the weighted mean as previously presented² and, for levofloxacin, was the adult clearance taken from Chien et al¹⁰. Only mean adult clearance values were regarded in this study.

Allometric clearance scaling

This approach to clearance scaling makes use of an allometric equation for the ratio of body weights to a specified power. The power function, for the purposes of scaling clearance is $\frac{3}{4}$ ^{7,11}. The clearance in a child (CL_{child}) is calculated from the clearance in the adult (CL_{adult} , both given in flow units, e.g. mL/min) using the equation;

$$CL_{\text{child}} = CL_{\text{adult}} \times \left(\frac{BW_{\text{child}}}{BW_{\text{adult}}} \right)^{\frac{3}{4}} \quad (1)$$

where BW_{child} is the body weight of the child and BW_{adult} is the body weight of the adult. The mean body weight of the paediatric population from each study was used in the allometric equation. If this value was not reported in the study, the mean weight value was taken for the mean age (or middle of the age range in some cases) using information from the International Commission on Radiological Protection (ICRP)¹². If the weight for a group of premature neonates was not reported (five of the 25 studies), the weight value was set at 2.5 kg. The adult body weight was set at 70 kg for all calculations.

Physiology-based clearance scaling

This approach uses information regarding clearance pathways in adults and scales them to children. The compound-specific proportion of total adult clearance is delineated into the compound-specific pathways of clearance. In the case of renal clearance, based on body weight and age of the child, glomerular filtration and tubular secretion are scaled to children based on a predictive equation generated retrospectively in children of different ages following administration of compounds undergoing renal elimination¹³, later slightly modified to account for even younger children². For hepatic enzymatic clearance, an adult intrinsic clearance (CL_{INT}) is calculated based on the well-stirred liver model;

$$CL_{INT} = CL_H \frac{Q_H}{Q_H - CL_H} \cdot \frac{1}{f_u} \quad (2)$$

where Q_H is the hepatic blood flow, CL_H is the plasma clearance, CL_{INT} is the intrinsic hepatic clearance per gram of liver weight and f_u is the fraction unbound in plasma. The generated adult intrinsic clearance value is then multiplied by an ontogeny factor that represents the activity of the specific enzyme in relation to the age of the child. This new child-scaled intrinsic clearance is used to generate an age-specific plasma clearance (CL_H) calculated from the re-arranged equation 2 using the age-specific body weight, liver weight, liver blood flow and predicted fraction unbound (scaled from adults based binding protein concentrations in plasma)¹⁴. The clearance pathways are summed to generate a total plasma clearance for the child. Detailed information on this approach, and the scaled clearances from the data set, is available in our previous study².

Approach comparison

The ratios of experimentally predicted to observed clearances (ml/min), using allometric and physiology-based scaling, were calculated and plotted against age for each compound. The compounds were grouped according to the primary process of clearance, which included the processes of renal, CYP3A4, CYP1A2 and UGT2B7 clearance. The ratio of predicted to observed clearances for the remaining multiple pathway clearance compounds were graphed together. The line of unity at which the predicted value was equal to the observed value was plotted together with the lines for where predicted clearance was either twice or half the observed clearance.

To determine the age range for which allometric scaling was appropriate, and to determine if this age range depended on the process of clearance, the clearance of the eight compounds that are cleared via one prominent pathway were calculated and compared to the observed clearance values. The ratios of predicted to observed clearances were visually examined for bias. A visual assessment was preferred over a numerical assessment of bias¹⁵, because the latter would be unfair. It is understood *a priori* that the allometric equation overestimates clearance in the young age groups^{5,11}, since this approach can not account for the enzymatic ontogeny and maturation process of the eliminating organ. Thus, the bias for the allometric equation would be heavily dependent on the number of data points under the age at which it can no longer reasonably predict clearance. Thus, the age at which the data points were equally above and below the line of unity was noted.

To compare the two models of scaling clearance, the seven evaluation compounds² were assessed. A visual assessment of the resulting graphs was focused on the bias of data points around the line of unity across the entire age range.

Results

Figure 1 summarises the results of both scaling approaches, divided by the process responsible (> 85%) for clearance. For each process; renal (Figures 1a and b), CYP3A4 (Figures 1c and d), CYP1A2 (Figures 1e and f) and UGT2B7 (Figures 1g and h), the clearance was scaled via the $\frac{3}{4}$ allometric power model (left side figures) and via the physiology-based approach (right side figures). These compounds were used to build the models used in the physiology-based approach² and thus cannot evaluate the model. Thus the figures representing the physiology-based clearance scaling are only shown here for reader interest. In the case of the allometric approach, it did not accurately predict clearance below a certain age in all cases as evidenced by the systematic accumulation of data points above the line of identity towards the younger ages.

In the case of renal elimination (Figure 1a), the allometric equation was two fold off the observed clearance value at a young age; however, the allometric equation consistently overestimated clearance until an age of approximately two years at which point allometric scaling consistently underestimated clearance.

There is evidence to suggest that for physiological processes an allometric power function of $\frac{2}{3}$ is more appropriate. Figure 2 presents the results using the allometric approach using a power function of $\frac{2}{3}$ in the allometric equation for gentamicin and isepamicin. In this case, the overestimation in children less than two years of age was greater than using the $\frac{3}{4}$ power function. For children over two years of age, there was consistent underestimation of clearance but with less bias than using the $\frac{3}{4}$ power function.

For the hepatic enzymes, the consistency of data above the line of identity appeared to be both age and process-dependent. The age at which the allometric equation produced little bias around the line of unity was within the first year for compounds eliminated primarily via CYP3A4 (Figure 1c), four years for CYP1A2 (Figure 1e) and six months for UGT2B7 (Figure 1g).

Figure 3 presents the results for which a comparison of the scaling approaches was made. Allometric scaling systematically overestimated clearance in the young age groups as evidenced




Figure 1 The predicted/observed clearance ratio plotted against age for compounds eliminated via renal clearance due to glomerular filtration (a, b), CYP3A4 (c, d), CYP1A2 (d, e) and UGT2B7 (f, g). The results using the $\frac{3}{4}$ allometric clearance scaling approach are shown on the left and the physiology-based approach results are in the right column. The solid line represents the line of identity and the dotted lines indicate where predicted clearance was twice or half the observed clearance. The arrow indicates data from premature neonates.




Figure 2 The predicted/observed clearance ratio plotted against age for gentamicin and isepamicin. Predicted clearances were derived using the $2/3$ allometric power function. The solid line represents the line of identity and the dotted lines indicate where predicted clearance was twice or half the observed clearance. The arrow indicates data from premature neonates.

by the lack of data points at or below the line of identity (Figure 3a). The physiology-based approach appeared not to be biased towards either over or under estimation of clearance across the age range of term neonates to 18 years (Figure 3b). Clearance in premature neonates was slightly overestimated (Figure 3b).

Discussion

The scaling of clearance from adults to children has been the subject of recent studies. There are two primary approaches proposed to achieve this; namely allometric scaling^{4-7,11,16} and physiology-based scaling^{2,5,17,18}. There is currently not enough data to suggest when the use of these scaling approaches is appropriate and/or feasible. An approach is required, or an integration of approaches, that can accurately predict the general trends associated with the age-dependence of clearance.

The $3/4$ power model has been used to scale metabolic and physiological processes among species based on body size, including scaling to humans (for a list of key references on the history of the $3/4$ power model see Anderson and Meakin⁷). This easy-to-use equation presents a significant improvement over other size models such as the surface area or per kg models¹¹. As expected, when the maturity of the processes responsible for clearance were immature, the allometric equation was not appropriate for predicting clearance, despite the inter-individual variability in the clearance values that are common in paediatric populations. Allometric scaling tends to systematically overestimate clearance in young children,

where the intrinsic activity of the eliminating process has not yet reached the activity in the adult. This was demonstrated elegantly in a study that compared the allometric approach to the physiology-based approach for the two CYP3A4 substrates, midazolam and alfentanil⁵.

In an effort to use the allometric approach in young children, others⁶ have examined the use of power functions other than $3/4$. As observed in our study, the $3/4$ function was appropriate in most cases for children over the age of five years. In younger children however, the power function that most accurately predicted clearance was age and drug dependent and no generalisations could be made as when these different power functions should be used⁶. This was likely due to the fact that the allometric approach takes into account size but not the ontogeny of the clearance pathway, which is both age (ontogeny) and drug (pathway-specific) dependent, as demonstrated in our study. The physiology-based approach, which explicitly accounts for the age dependence of enzymatic activity, predicted the trends of clearance in children.

As discussed in the Bjorkman study⁵ for CYP3A4, the *in vitro* assays used to determine the age-dependence of enzyme activity can vary greatly depending on the substrate used to generate the data. In his study, using these different *in vitro* ontogeny values led to greatly varying clearance predictions for children. In our previous study², both literature based clearance values for midazolam and alfentanil and *in vitro* activity data were used to build the ontogeny model of CYP3A4 for children, thus not relying solely on




Figure 3 The predicted/observed clearance ratio plotted against age for compounds eliminated via multiple pathways. Graph a is based on clearance scaling using the $\frac{3}{4}$ power allometric approach. Graph b is based on the physiology-based clearance scaling approach. The solid line represents the line of identity and the dotted lines indicate where predicted clearance was twice or half the observed clearance. The arrow indicates data from premature neonates.

the variable *in vitro* data. This was later used to predict the hepatic clearance of other CYP3A4 substrates in children with a good agreement². Using the latter approach to physiology-based clearance scaling, the accuracy of clearance predictions was independent of the age with the exception of premature neonates, which tended to be slightly overestimated.

Premature neonates represent a specific subgroup, with variable clearances dependent on weight, gestational and postnatal age. For example, to define caffeine clearance following intravenous administration in premature neonates, Lee et al¹⁹ had to generate an empirically-based equation that included both current weight and postnatal age, with postnatal age being the more important of the two. Furthermore, premature neonates generally receive drugs for varying clinical conditions, which may alter the neonate's physiology and hence the drug's pharmacokinetics. Using the physiology-based approach, all premature experimental groups were considered equivalent such that all premature neonates in a compound group were given the same clearance value. Clearance pathway enzymatic ontogeny was considered in the physiology-based approach. Therefore, this more accurately predicted premature clearances in comparison to the allometric approach.

In the case of passive renal elimination via glomerular filtration, it appeared that the allometric equation did not accurately scale clearance. Since the $\frac{3}{4}$ power model has primarily been used to scale active physiological and metabolic processes, the inability to accurately scale the passive process of glomerular filtration may have been expected. For passive processes, an allometric power function of $\frac{2}{3}$ may be more appropriate as this is directly related to body surface area. While the underestimation in children over two years of age was less than using the $\frac{3}{4}$ power function, the overestimation in children under two years of age was greater. For renal clearance, other means of scaling should be used^{2,13}.

Interestingly, it appeared that at the age below which the allometric equation was no longer accurate, closely matched the age at which the enzymes responsible reached intrinsic adult activities. For example, the allometric equation seems appropriate for scaling the clearance of the UGT2B7 and CYP3A4 compounds in children over the age of at least six months. This corresponds to the age at which *in vitro* enzyme activity reaches adult activity for UGT2B7³ and CYP3A4³. CYP1A2 is the last major enzyme to develop reaching adult activity levels well beyond one year of age²⁰, up to potentially 15 years of age³. The age at which the

allometric equation correctly predicted clearance of the compounds primarily eliminated via this enzyme was approximately four years. More confidence could be gained in determining the youngest age at which the allometric approach is appropriate for scaling clearance with an increase in the quite sparse data for neonates between birth and one year of age. Unfortunately, a complete range of ages was not available for all compounds potentially skewing these results. Regardless, it appears from this study that the age at which the allometric equation becomes a useful means of scaling clearance is following the age at which the maturity of the process is complete or near to complete and, is thus, process specific.

These studies represent those of normal weight for age children, based on a comparison of the reported average weights to standard weights in children from the ICRP publication¹². We did not have enough data to support the assertion that the allometric equation could be used to predict clearance in overweight and underweight children. Holford¹¹ suggests that in cases of unusual body composition, a good place to start is using size based on the normal weight for age. Only with individual clearance data from children with unusual body composition can we begin to test the sensitivity of clearance scaling to altered body composition.

This study has shown that the appropriate use of the allometric equation depends on two factors; the age of the child and the primary process responsible for clearance. While this study only focused on four primary processes of clearance, renal, CYP3A4, CYP1A2 and UGT2B7, the age-dependent predictivity of the allometric equation varied greatly. As has been demonstrated previously², a physiology-based approach is predictive of these processes from birth (term) to 18 years, however; the physiology-based approach requires more input and is mathematically more challenging. Because of the ease of use of the allometric equation, it is suggested to have an integrative approach that uses the allometric approach when appropriate and switches to the physiology-based approach in the younger age range. An advantage of the physiology-based approach is that clearance in a child due to a medical condition, such as a reduced eliminating organ blood flow, can be assessed with greater confidence than using a body weight scaling approach.

Clearance scaling is a first step towards the scaling of pharmacokinetic profiles from adults to children. The next step is a determination of the age-dependence of distribution volumes using either an allometric approach⁷ or a physiological approach⁹. Together, these two parameters

can lead to a reasonable prediction of a child's pharmacokinetic profile and decisions regarding dosing and potential therapeutic and/or adverse effect events can be informed.

Competing interests

None declared

References


1. Choonara I. Unlicensed and off-label drug use in children: implications for safety. *Expert Opin Drug Saf* 2004;3:81-83.
2. Edginton AN, Schmitt W, Voith B et al. A mechanistic approach for the scaling of clearance in children. *Clin Pharmacokinet* 2006;45:683-704.
3. Alcorn J, McNamara PJ. Ontogeny of hepatic and renal systemic clearance pathways in infants: part I. *Clin Pharmacokinet* 2002;41:959-998.
4. Knibbe CA, Zuideveld KP, Aarts LP et al. Allometric relationships between the pharmacokinetics of propofol in rats, children and adults. *Br J Clin Pharmacol* 2005;59:705-711.
5. Bjorkman S. Prediction of cytochrome P450-mediated hepatic drug clearance in neonates, infants and children: how accurate are available scaling methods? *Clin Pharmacokinet* 2006;45:1-11.
6. Mahmood I. Prediction of drug clearance in children from adults: a comparison of several allometric methods. *Br J Clin Pharmacol*. 2006;61:545-557.
7. Anderson BJ, Meakin GH. Scaling for size: some implications for paediatric anaesthesia dosing. *Paediatr Anaesth* 2002;12:205-219.
8. Van der Marel CD, Anderson BJ, Van Lingen RA et al. Paracetamol and metabolite pharmacokinetics in infants. *Eur J Clin Pharmacol* 2003;59:243-251.
9. Edginton AN, Willmann S, Schmitt W. Development and evaluation of a generic physiology-based pharmacokinetic (PBPK) model for children. *Clin Pharmacokinet* 2006;45:1013-1034.
10. Chien S, Wells TG, Blumer JL et al. Levofloxacin pharmacokinetics in children. *J Clin Pharmacol* 2005;45:153-160.
11. Holford NH. A size standard for pharmacokinetics. *Clin Pharmacokinet* 1996;30:329-332.
12. International Commission on Radiological Protection (ICRP). Basic anatomical and physiological data for use in radiological protection: reference values. Publication 98. Amsterdam, The Netherlands: Elsevier Science, 2002.
13. Hayton WL. Maturation and growth of renal function: dosing renally cleared drugs. *AAPS PharmSci* 2002; 2:1-7.
14. McNamara PJ, Alcorn J. Protein binding predictions in infants. *AAPS Pharm Sci* 2002;4:1-8.
15. Sheiner LB, Beal SL. Some suggestions for measuring predictive performance. *J Pharmacokinet Biopharm* 1981;9:503-512.
16. Anderson BJ, Pons G, utret-Leca E et al. Pediatric intravenous paracetamol (propacetamol) pharmacokinetics: a population analysis. *Paediatr Anaesth* 2005; 5:282-292.
17. Alcorn J, McNamara PJ. Ontogeny of hepatic and renal systemic clearance pathways in infants: part II. *Clin Pharmacokinet* 2002;41:1077-1081.
18. Bjorkman S. Prediction of drug disposition in infants and children by means of physiologically based pharmacokinetic (PBPK) modelling: theophylline and midazolam as model drugs. *Br J Clin Pharmacol* 2004;59:691-704.
19. Lee TC, Charles B, Steer P et al. Population pharmacokinetics of intravenous caffeine in neonates with apnea of prematurity. *Clin Pharmacol Ther* 1997;61:628-640.
20. Sonnier M, Cresteil T. Delayed ontogenesis of CYP1A2 in the human liver. *Eur J Biochem* 1998;251:893-898.

CrossRefs are available in the online published version of this paper:
<http://www.librapharm.com>
 Paper PPDT-0167_5, Accepted for publication: 11 October 2006
 Published Online: 15 November 2006
 doi:10.1185/146300906X148530

Paediatric and Perinatal Drug Therapy

Instructions to Authors

1. All manuscripts should be in the English language. They should be submitted to Imti Choonara, Academic Division of Child Health, (University of Nottingham) The Medical School, Derbyshire Children's Hospital, Uttoxeter Road, Derby DE22 3DT, UK (Email: imti.choonara@nottingham.ac.uk). Manuscripts from North America should be submitted to Professor Greg Kearns, Children's Mercy Hospital, 2401 Gilham Road, Kansas City, Missouri 64108, USA (Email: gkearns@cmh.edu). *Paediatric and Perinatal Drug Therapy* is published, produced and distributed by Informa Healthcare, Telephone House, 69-77 Paul Street, London EC2A 4LQ, UK. Tel: +44 (0) 20 7017 7665; Fax: +44 (0) 20 7017 7831.
2. Electronic submission of manuscripts facilitates rapid publication. Most common word-processor formats are acceptable, although Microsoft Word is preferred. If electronic submission is not possible, then a single copy of the manuscript with an accompanying disk is acceptable. The manuscript should be double-spaced and on one side of the paper only. Each paper should contain the following: (a) a short descriptive title, (b) the name(s) and initials of the author(s), (c) the Centre at which the work was carried out or the location of the author(s), (d) a summary or abstract of the main facts and results (e), an Introduction, (f) separate main sections, (g) a final Discussion or Conclusions section, (h) any acknowledgements and (i) full references to relevant material in the text. Authors are also requested to supply approximately six 'key words', in English, preferably from the Index Medicus Medical Subject Heading (MeSH) list.
3. All drugs and other compounds should be referred to by their internationally accepted generic names and not by individual company trade marks, unless it is essential for clarity, as in the case of combination products, or to avoid confusion, e.g. between different formulations. Specialised abbreviations and symbols should not be used unless first explained in the text. Dosages and measurements should be given in the units in which they were made, but non-metric units should be accompanied by metric (SI) equivalents.
4. Acknowledgement must be given by authors of grants, fellowships, or any commercial assistance received or of any affiliation which is relevant to the work reported.
5. All references should be individually numbered in Arabic numerals and cited where they appear in the text. At the end of the paper, references should be listed in strict numerical order. The names of all authors for each reference must be given (unless there are six or more, in which case the first three should be listed, followed by 'et al.'). They should be followed by: (a) the full title of the paper, (b) the abbreviated title of the journal (ANSI/BSI system), (c) the year of publication, and (d) the volume and page number(s). Reference to books must give the publisher, place and year of publication, name(s) of the editor(s) where authorship is multiple, and first page number of chapter referred to.
6. All tables and illustrations should be provided with short descriptive legends, numbered consecutively, and their relevant position in the text clearly indicated. Tables should have concise headings to all columns and be identified by Arabic numerals, e.g. Table 2. They should be supplied within the files on disk in cellular form rather than in simple tabbed form. Line diagrams should be supplied both as files on disk in either .TIF or .EPS format and in the format of the program used to produce them. If this is not possible, they should be supplied in a suitable finished form for reproduction and in proportion to the single-column width (80 mm) or double-column width (165 mm). Photographic illustrations will usually be accepted. Illustrations should also be identified by Arabic numerals, e.g. Figure 2.
7. Papers are published on the understanding that their copyright becomes the property of the Publishers once they are accepted for publication. Authors must state clearly if the paper is being actively considered for publication or has been published elsewhere in the world. If subject to copyright (and this includes illustrations), copyright clearance is the sole responsibility of the author and must be supplied in writing to the Publishers. Papers first published in *Paediatric and Perinatal Drug Therapy* must not be translated, abridged or reprinted in any form elsewhere in the world without the written consent of the Publishers.
8. Proofs in page form will be sent to the main author for checking provided that this will not result in delayed publication of any issue of the journal. If, because of postal delays, etc. time is limited, the Publishers reserve the right to have proofs checked against original manuscripts by their editorial staff and/or the editors. No major alterations to text will be accepted at proof stage.



PAEDIATRIC CLINICAL PHARMACOLOGY

Edited by

Evelyne Jacqz-Aigrain *Hopital Robert Debre, Paris, France*

Imti Choonara *University of Nottingham, UK*

January 2006 / ISBN: 0824721896 / 500 pp. / List Price: \$169.95

Special Offer: \$152.95 (Please refer to discount code DFL05DK20)

The treatment of children with medicinal products is an important scientific area. It is recognised that many medicines used extensively in paediatric patients are either unlicensed or off-label. This textbook will help paediatric health professionals effectively treat children with the most appropriate medicine with minimal side effects.

This source covers

- the generic principles behind the use of medicines in children
- the science of paediatric clinical pharmacology, clinical trials, and aspects relating to formulation development
- specific clinical areas encompassing the fetus, neonate, and child

Contents: Drug Evaluation in Children • Paediatric Clinical Pharmacology • The Administration of Medicines to Children • Drugs in Pregnancy and Lactation • Medicines in Neonates • Infections • Critical Care, Neurology, and Analgesia • Immunosuppressants, Rheumatic and Gastrointestinal Topics: Clinical Section (A) • Respiratory, Endocrine, Cardiac, and Renal Topics: Clinical Section (B)

To order, contact your nearest Informa Healthcare/Taylor and Francis office:

North & South America:

Informa Healthcare, Kentucky Distribution, 7625 Empire Drive, Florence, KY 41042

Tel: (800) 634-7064

Fax: (800) 248-4724

e-mail: orders@taylorandfrancis.com

UK and International (excluding East & Southeast Asia)

Taylor & Francis, Customer Services, TPS, Cheriton House, North Way, Andover, Hampshire SP10 5BE, UK

Tel (UK): +44 (0) 1264 34 2926

Tel (Int'l): +44 (0) 1264 34 3070

Fax: +44 (0) 1264 34 3005

e-mail (UK): uk.tandf@thomsonpublishingservices.co.uk

e-mail (Int'l): international.tandf@thomsonpublishingservices.co.uk

East and Southeast Asia

Singapore Sales Office, Taylor & Francis Asia Pacific, 240 Macpherson Road, #08-01 Pines Industrial Building, Singapore 348574

Tel: +65 6741 5166

Fax: +65 6742 9356

e-mail: sales@tandf.com.sg

ESDP–Eudipharm

5th European Course

Evaluation of Medicinal Products in Children

Part I

22–24 February 2007, Paris

- Specific aspects of paediatric pharmacology
- Epidemiology of drug use in children
- Legislation and licensing
- Clinical trials in children
- Pharmacogenomics
- Ethical issues
- Drug development
- Post-marketing surveillance in children

*For further details, please email:
gerard.pons@svp.ap-hop-paris.fr*

ESDP–Eudipharm

5th European Course

Evaluation of Medicinal Products in Children

Part II

14–17 March 2007, Paris

- Pre/perinatal drug evaluation and use
- Placental drug transfer
- Drug evaluation and protocol design
in various therapeutic areas in children
(workshop format)
 - oncology
 - epilepsy
 - pain
 - endocrinology
 - neonatology

*For further details, please email:
gerard.pons@svp.ap-hop-paris.fr*

Introduction to Paediatric and Perinatal Drug Therapy

Edited by I Choonara, A J Nunn and G Kearns

The aim of this book is to improve the use of drug therapy in paediatric patients of all ages. Clear and concise chapters offer an insight into key aspects of paediatric clinical pharmacology and pharmacy.

- Introduction to clinical pharmacokinetics
- The effect of age on drug metabolism
- Drug toxicity
- Ethical issues in research with children
- Licensing
- Pharmaceutical industry viewpoint
- Extemporaneous formulations
- Advances in oral formulation development
- Inhalational devices for young children
- Placental transfer of drugs
- Surfactant therapy in neonates
- The use of aminoglycosides in neonates
- Antibiotics for acute otitis media in infancy
- Pain assessment in children
- Medication errors

Paperback 238 pages

ISBN 1-897676-37-9

£25.00 (approx. US\$46.00 or 36.00 Euros)

To order your copy, please contact: Nottingham University Press,
Manor Farm, Church Lane, Thrumpton, Nottingham NG11 0AX, UK

Tel: +44 (0) 115-983 1011 Fax: +44 (0) 115-983 1003

Email: orders@nup.com

Make better research decisions, faster, with an...



Expert Opinion

Ten phase-specific, peer-reviewed journals for pharmaceutical R&D,
from drug discovery to pharmacotherapy

Go online today to learn more about the series, request your
FREE TRIAL and sign up for your table of content e-alerts too!

Ref: JEO 0020A

www.expertopin.com

informa
healthcare

