

Indomethacin Treatment of Patent Ductus Arteriosus in Premature Infants

John McIntyre Senior Lecturer in Child Health

Academic Division of Child Health (University of Nottingham), Derbyshire Children's Hospital, Uttoxeter Road, Derby DE22 3NE, UK. email: john.mcintyre@nottingham.ac.uk

Bart Van Overmeire Senior Lecturer in Neonatology

Department of Paediatrics, University Hospital of Antwerp, Wilrijkstreet 10, B-2650 Edegem, Belgium

John N. van den Anker Professor of Paediatrics and Neonatology

Sophia Children's Hospital, Rotterdam, The Netherlands

Abstract

Shortly after birth, physiological processes usually effect prompt functional closure of the ductus arteriosus. With increasing prematurity there is an increased likelihood of the ductus arteriosus remaining patent. To overcome the undesirable effects of left to right shunting of blood through the patent ductus arteriosus (PDA), closure by surgical ligation and medical treatment have become established practice. Since the mid-1970s, indomethacin, an inhibitor of prostaglandin synthesis, has been used to promote closure of a PDA. Despite longstanding use and numerous clinical trials, the optimal management remains unclear. This overview covers the pharmacology of indomethacin in the neonate and its use for treating PDA in the premature infant.

Key words: Indomethacin – Ductus arteriosus, patent – Infant, premature

Introduction

In fetal circulation the ductus arteriosus channels blood from the pulmonary artery directly into the descending aorta. This enables about 90% of right ventricular output to bypass the unexpanded lungs. At birth, functional closure of this channel redirects blood flow to the pulmonary vascular bed and allows the lungs to start effective respiratory gas exchange. Closure involves interaction of oxygen, prostaglandins, nitric oxide and the unique musculature of the ductus arteriosus¹⁻³. In healthy term babies, ducts are functionally closed in 20% on day 1, 82% by day 2, 96% by day 3 and 100% by day 4⁴.

These mechanisms do not always occur promptly in premature infants. A patent ductus arteriosus

(PDA) is found in about 40% of infants of < 1000 g birthweight on day 3⁵, and, in infants of < 28 weeks' gestation, spontaneous closure by day 5 occurs in only 27%⁶. Left to right shunting of blood through a haemodynamically significant PDA increases pulmonary blood flow and cardiac workload, and reduces renal and gastrointestinal blood flow. A PDA has been thought to increase the severity of respiratory distress^{7,8} and the risk of bronchopulmonary dysplasia⁹.

Surgical ligation and medical drugs, most commonly indomethacin, are used to treat infants with a haemodynamically significant PDA. Pharmacological closure of a PDA in premature infants was first reported in 1976^{10,11} and subsequent studies show it to be effective in about 80% of treated infants¹². Indomethacin has

also been used to prevent a symptomatic PDA developing in at-risk infants¹³. However, despite over 30 years of various approaches, the optimum regimen for the management of PDA in prematurity has still not been unequivocally established.

Pharmacology of Indomethacin

Indomethacin is a methylated indole derivative and a non-selective potent inhibitor of the prostaglandin-forming cyclo-oxygenase. It is highly bound to serum proteins. Metabolism via *O*-methylation, deacylation and conjugation to inactive metabolites allows for elimination via the biliary route; < 10% is excreted unchanged in the urine³.

Intravenous indomethacin in adults has a plasma half-life of 2.6–11.2 h and a plasma clearance of 44–109 ml/kg/h^{14,15}. Defining pharmacokinetic parameters in the neonate is hampered by the many physiological changes occurring after birth. Despite their limitations, pharmacokinetic studies that have been attempted do give some insight. With oral administration, absorption of indomethacin is incomplete and variable¹⁶. For intravenous indomethacin, studies have used different doses (0.2, 0.25 and 0.3 mg/kg) but all show large inter-individual variations and important differences compared with the adult data: plasma half-life ($t_{1/2}$: 11–36 h) is much longer and the clearance rate (7–16 ml/kg/h) much less^{17–20}. There are also large inter-individual variations in the apparent volume of distribution, and the duct being open or closed may in part contribute to this¹⁹. Increasing postnatal age, not gestational age, is an important determinant of increasing indomethacin clearance and volume of distribution. A study of the population pharmacokinetic parameters of indomethacin found similar results²¹.

The relationship between the plasma concentration of indomethacin and the desired effect of ductal closure is controversial. In some studies plasma concentration has not correlated with efficacy^{12,22,23}. In contrast, others have found a relationship^{19,21,24–28}. Various effective concentrations are suggested: 1 mg/l 10 h post-dose²⁴; 0.5 mg/l 8 h post-dose²⁵; and 0.25 mg/l over a three-day course²⁷. From population pharmacokinetic data, the mean serum concentration at PDA closure was 1.43 mg/l²¹. Furthermore, low plasma levels of indomethacin, short $t_{1/2}$ and more efficient clearance, are associated with unsuccessful indomethacin infusions¹⁹.

Surgical Treatment

From the early 1970s surgical ligation has been used in premature neonates with a PDA, but, with increasing use of indomethacin surgery, is more often reserved for infants with contraindications to, or failure of, medical treatment. The National Collaborative Study suggested surgery as a backup treatment¹². However, some reports recommend surgery as the first-line approach in extremely low-birth-weight infants. The perioperative complications associated with surgery include hypertension, pneumothorax, chylothorax, ductal tear, minor wound infections and vocal cord paralysis. Morbidity rates reported are 1–16% and mortality 0–10%^{29–32}. In experienced centres with low morbidity and mortality, surgery may be an appropriate alternative in extremely low-birth-weight infants where indomethacin has a high failure rate and significant adverse effects.

Medical Treatment

Timing of Treatment

With respect to timing, medical treatment may be prophylactic (usually within 24 hours of birth), early (2–3 days) or late (8–10 days). Clyman³³ identified four general treatment strategies that have been studied with respect to timing:

- Late symptomatic vs. no backup.
- Late symptomatic vs. later backup.
- Early symptomatic vs. late symptomatic.
- Prophylactic vs. early symptomatic.

The only study of Strategy 1 was too small to draw firm conclusions, but ductal closure did appear associated with reduced mechanical ventilation, necrotising enterocolitis (NEC) and retinopathy of prematurity (ROP)³⁴. In the other studies, a backup treatment (indomethacin or ligation) was available. From meta-analysis of the studies available in 1996³³, the group receiving early symptomatic treatment with indomethacin had a reduction in bronchopulmonary dysplasia (OR = 0.39, CI 0.21–0.76; $p < 0.005$), need for mechanical ventilation ($p < 0.005$) and NEC (OR = 0.24; CI = 0.06–0.96; $p < 0.05$) compared with late symptomatic treatment. Prophylactic treatment did not have any advantage in long-term pulmonary outcomes or NEC compared with early symptomatic treatment; however, the incidence of grade 3 and 4 intraventricular haemorrhage (IVH) was reduced (OR = 0.51; CI = 0.28–0.95; $p < 0.05$). In another systematic review and meta-analysis, Fowlie³⁵ found evidence that prophylactic indomethacin reduced the incidence of symptomatic PDA (estimate of risk difference –0.217; CI –0.275, –0.160) and the incidence of grades 3 and 4 IVH (estimate of risk difference

–0.039; CI –0.066, –0.011). However, mortality and other morbidity was not significantly affected.

The benefit of prophylactic indomethacin in reducing the incidence of IVH may be independent of its effect on a PDA. There is evidence of low-dose prophylactic indomethacin reducing IVH³⁶. More recently it has been suggested that prophylaxis may also reduce the incidence of periventricular leukomalacia (PVL)⁶.

Length of Treatment

A commonly used course of indomethacin is three doses. The National Collaborative Study used three doses at 12-hourly intervals: 0.2 mg/kg for infants in the first 2–7 days and 0.2 mg/kg followed by 0.25 mg/kg for the second and third dose for infants > 8 days. The initial response rate of duct closure (about 80%) and subsequent reopening in responders (about 30%) was broadly in keeping with the many previous studies¹². To improve sustained closure rates, prolonged courses of indomethacin have been suggested as an alternative^{24,37,38}.

Using indomethacin level monitoring to individually tailor doses it was suggested that maintaining a plasma level of 0.5 mg/l for a week would be effective in maintaining duct closure²⁴. In a study of 39 low-birth-weight babies with a PDA, after three doses of indomethacin at 0.2 mg/kg, babies were randomised to placebo or 0.2 mg/kg/day for five days³⁷. The maintenance regimen reduced the PDA recurrence rate and the need for surgical ligation. In a study of 121 babies, where PDA detection and follow-up was clinical, 0.1 mg/kg/day for six days was compared with three doses of 0.2 mg/kg 12-hourly³⁸. For the prolonged course, the initial response rate was better (90% vs. 77%) and of the responders the relapse rate (21% vs. 40%) was lower.

Other studies have not confirmed the advantages of a prolonged course. In a study of 70 low-birth-weight infants, randomisation was to just two doses of 0.15 mg/kg 12 hours apart or a maintenance regimen of 0.1 mg/kg/day for five days after the initial doses³⁹. The initial closure rate appeared better in the maintenance group but overall the incidence of reopening was not different. More recently short versus prolonged indomethacin therapy was compared in 61 preterm infants with a PDA, confirmed on echocardiography⁴⁰. Those receiving a short course had a shorter duration of oxygen supplementation and less frequent symptoms of NEC, but sustained closure rates (74% vs. 60%), mortality and other neonatal morbidity rates were similar. It was concluded that a prolonged course offered no advantages over a standard course in preterm

Table 1. Indomethacin for closure of PDA-licensed recommendations (from ABPI Compendium of Data Sheets and Summaries of Product Characteristics 1999–2000)

Age at first dose	Intravenous dose (microgram/kg)		
	1st dose	2nd dose	3rd dose
< 48 hours	200	100	100
2–7 days	200	200	200
> 7 days	200	250	250

infants with a haemodynamically significant PDA.

In the absence of a clear benefit for prolonged courses, the current licensed therapy (shown in Table 1) is three intravenous doses given at 12–24-hour intervals.

Other Effects of Indomethacin

As a non-specific prostaglandin synthesis inhibitor, indomethacin would be expected to have a spectrum of systemic effects, and these have been confirmed in clinical studies.

Cerebral

Doppler ultrasound and near infrared spectroscopy (NIRS) have been used to study the effects of indomethacin on cerebral haemodynamics. Doppler ultrasound shows significant reductions in cerebral blood flow velocities of 20–40% within minutes of the injection that persist for one to two hours^{41–44}. However, infusion over 20 minutes did not result in significant changes⁴³ and a slow continuous infusion effectively closed a PDA without decreasing cerebral blood flow velocities⁴⁵. NIRS shows substantial reductions in cerebral blood flow, cerebral oxygen delivery, cerebral blood volume and cerebral blood volume reactivity^{46–48}. The underlying mechanism of these changes is unclear and they are not seen with another prostaglandin inhibitor, ibuprofen^{47,48}, raising the possibility of a mechanism other than prostaglandin synthesis inhibition. The alteration of cerebral haemodynamics in infants already vulnerable to cerebral hypoperfusion remains an area of concern.

There is evidence that early use of indomethacin may protect against IVH^{36,49,50} and PVL⁶. Meta-analysis of trials using indomethacin prophylaxis³⁵ suggests the incidence of severe IVH is reduced by about 40% (estimate of risk difference –0.039; CI –0.066, –0.011). Furthermore, data from the Multicentre Randomised Indomethacin IVH Prevention Trial showed no adverse neuro-

developmental effects at 4.5 years in those who had received early low-dose indomethacin compared with those who had received placebo⁵¹. In this study, the cerebral palsy rate was the same in both groups (7%) but the proportion of children with no, or minor, developmental handicap was smaller in the indomethacin group.

Renal

Both a PDA and indomethacin treatment can alter renal function³. Indomethacin can result in oliguria, increased serum blood urea nitrogen and creatinine, decreased glomerular filtration rate and serum electrolyte imbalance. The renal dysfunction associated with indomethacin can in part be explained by a reduction in renal blood flow and perfusion^{52–53}. Associated changes also include a decrease in plasma renin activity and a rise in arginine vasopressin⁵⁴.

These effects are usually transitory without resulting in prolonged renal dysfunction⁵⁵. Infants receiving frusemide with indomethacin have higher urine output, glomerular filtration rates and fractional excretion of sodium⁵⁶. However, studies of frusemide in indomethacin-treated infants have been small with methodological limitations. Meta-analysis of these highlighted dehydration as a contraindication to frusemide use and concluded that the risk–benefit ratio could only be assessed by a large randomised trial⁵⁷. Dopamine does not reduce the renal side-effects of indomethacin⁵⁸.

Gastrointestinal

Bowel perforation and necrotising enterocolitis are reported in infants with a PDA who are treated with indomethacin⁵⁹; isolated perforation without NEC is also well described⁶⁰.

Doppler studies show a pronounced and sustained fall in the velocity of the superior mesenteric artery shortly after administration of indomethacin^{61,62}; this is less and later if the indomethacin is given by infusion rather than as a bolus⁶². The further reduction in flow to an already compromised vascular bed helps explain some of the observed gastrointestinal effects. The mechanism remains unclear and it has been suggested may be independent of its effect on prostaglandins⁶².

Despite the potential for adverse effects on the GI tract, early use of indomethacin may reduce the incidence of NEC overall³³. Meta-analysis of studies of prophylactic indomethacin found the effect of indomethacin on the incidence of NEC did not reach statistical significance, although there is, perhaps, a trend towards increased risk¹³.

Haematological

Indomethacin may interfere with platelet function and result in impaired haemostasis⁶³. Although reduced platelet counts after treatment are reported with prophylactic indomethacin³⁶, there does not appear to be a significant difference between treatment and placebo groups³⁵.

Alternative Prostaglandin Inhibitors

Because of the side-effects of indomethacin, alternative prostaglandin inhibitors have been tried: aspirin, mefenamic acid, ethamsylate, sulidnac and ibuprofen^{10,64–68}. Ibuprofen appears to offer effective closure of the PDA without notable early adverse effects^{67,68}. Furthermore, it seems to have less effect on cerebral, mesenteric and renal haemodynamics^{48,69}. A recent study of ibuprofen used on the third day of life concluded it was as efficacious as indomethacin for treatment of a PDA and less likely to induce oliguria⁷⁰. Long-term follow-up studies will be required to determine the precise role of ibuprofen in treating PDA.

Conclusion

For over 30 years closure of the ductus arteriosus in premature infants has been pursued by surgical and pharmacological means. From this accumulated experience, what conclusions can be drawn?

A PDA is common in premature infants with RDS and is present in 40% of infants < 1000 g on day 3. Persistent large left to right shunting is undesirable. Interventions to promote PDA closure need to be carefully considered as it will close spontaneously in many infants. Pharmacotherapy is the first-line approach in most cases, but for very low-birth-weight infants, surgery is predictable, and, in experienced centres, has low morbidity and mortality^{29–32}. Indomethacin is effective and the most commonly used drug treatment, but controversy still surrounds optimum timing and dosing strategies. Early treatment in the first 2–4 days, before clinical evidence of a large shunt with congestive heart failure develops, reduces pulmonary morbidity and NEC compared with late (8–10 days) treatment³³. Prophylactic indomethacin exposes all infants to its wide range of potential adverse effects and does not appear to have benefit in terms of mortality and pulmonary morbidity compared with early symptomatic treatment^{33,35}. However, prophylaxis does result in a higher rate of permanent ductal closure⁶, appears to have a benefit of protecting against severe IVH and, at

present, there is no long-term evidence of harm^{13,51}. The standard course is three doses at 12-hour intervals, but evidence is conflicting on whether a prolonged course has additional benefit. Pharmacological closure of a PDA without side-effects remains a challenge, although recent studies of ibuprofen^{47,48,67-70} show some promising new developments.

References

- Clyman RI, Chan CY, Mauray F, et al. Permanent anatomic closure of the ductus arteriosus in newborn baboons: the roles of postnatal constriction, hypoxia, and gestation. *Pediatr Res* 1999;45:19-29
- Clyman RI, Waleh N, Black SM, Riemer RK, Mauray F, Chen YQ. Regulation of ductus arteriosus patency by nitric oxide in fetal lambs: the role of gestation, oxygen tension, and vasa vasorum. *Pediatr Res* 1998;43:633-44
- Hammerman C. Patent ductus arteriosus. Clinical relevance of prostaglandins and prostaglandin inhibitors in PDA pathophysiology and treatment. *Clin Perinatol* 1995;22:457-79
- Lim MK, Hanretty K, Houston AB, Lilley S, Murtagh EP. Intermittent ductal patency in healthy newborn infants: demonstration by colour Doppler flow mapping. *Arch Dis Child* 1992;67:1217-8
- Ellison RC, Peckham GJ, Lang P, et al. Evaluation of the preterm infant for patent ductus arteriosus. *Pediatrics* 1983;71:364-72
- Narayanan M, Cooper B, Weiss H, Clyman RI. Prophylactic indomethacin: factors determining permanent ductus arteriosus closure. *J Pediatr* 2000;136:330-7
- Thibeault DW, Emmanouilides GC, Nelson RJ, Lachman RS, Rosengart RM, Oh W. Patent ductus arteriosus complicating the respiratory distress syndrome in preterm infants. *J Pediatr* 1975;86:120-6
- Siassi B, Blanco C, Cabal LA, Coran AG. Incidence and clinical features of patent ductus arteriosus in low-birthweight infants: a prospective analysis of 150 consecutively born infants. *Pediatrics* 1976;57:347-51
- Brown ER. Increased risk of bronchopulmonary dysplasia in infants with patent ductus arteriosus. *J Pediatr* 1979;95(5 Pt 2):865-6
- Heymann MA, Rudolph AM, Silverman NH. Closure of the ductus arteriosus in premature infants by inhibition of prostaglandin synthesis. *N Engl J Med* 1976;295:530-3
- Friedman WF, Hirschklau MJ, Printz MP, Pitlick PT, Kirkpatrick SE. Pharmacologic closure of patent ductus arteriosus in the premature infant. *N Engl J Med* 1976;295:526-9
- Gersony WM, Peckham GJ, Ellison RC, Miettinen OS, Nadas AS. Effects of indomethacin in premature infants with patent ductus arteriosus: results of a national collaborative study. *J Pediatr* 1983;102:895-906
- Fowle PW. *Cochrane Database of Systematic Reviews* 2000:1
- Alvan G, Orme M, Bertilsson L, Ekstrand R, Palmer L. Pharmacokinetics of indomethacin. *Clin Pharm Ther* 1975;18:364-73
- Kwan KC, Breault GO, Umbenhauer ER, McMahon FG, Duggan DE. Kinetics of indomethacin absorption, elimination, and enterohepatic circulation in man. *J Pharm Biopharm* 1976;4:255-80
- Evans MA, Bhat R, Vidyasagar D, Vadapalli M, Fisher E, Hastreiter A. Gestational age and indomethacin elimination in the neonate. *Clin Pharm Ther* 1979;26:746-51
- Thalji AA, Carr I, Yeh TE, Raval D, Luken JA, Pildes RS. Pharmacokinetics of intravenously administered indomethacin in premature infants. *J Pediatr* 1980;97:995-1000
- Yaffe SJ, Friedman WF, Rogers D, Lang P, Ragni M, Saccar C. The disposition of indomethacin in preterm babies. *J Pediatr* 1980;97:1001-6
- Brash AR, Hickey DE, Graham TP, Stahlman MT, Oates JA, Cotton RB. Pharmacokinetics of indomethacin in the neonate. Relation of plasma indomethacin levels to response of the ductus arteriosus. *N Engl J Med* 1981;305:67-72
- Gal P, Ransom JL, Weaver RL, et al. Indomethacin pharmacokinetics in neonates: the value of volume of distribution as a marker of permanent patent ductus arteriosus closure. *Ther Drug Monitor* 1991;13:42-5
- Wiest DB, Pinson JB, Gal PS, et al. Population pharmacokinetics of intravenous indomethacin in neonates with symptomatic patent ductus arteriosus. *Clin Pharm Ther* 1991;49:550-7
- Smith IJ, Goss I, Congdon PJ. Intravenous indomethacin for patent ductus arteriosus. *Arch Dis Child* 1984;59:537-41
- Ramsay JM, Murphy DJ, Jr, Vick GWD, Courtney JT, Garcia-Prats JA, Huhta JC. Response of the patent ductus arteriosus to indomethacin treatment. *Am J Dis Child* 1987;141:294-7
- Seyberth HW, Knapp G, Wolf D, Ulmer HE. Introduction of plasma indomethacin level monitoring and evaluation of an effective threshold level in very low birth weight infants with symptomatic patent ductus arteriosus. *Eur J Pediatr* 1983;141:71-6
- Lewis IG, Harvey DP, Maxwell GM. Indomethacin therapy of patent ductus in preterm infants controlled by plasma levels. *Austra Paed J* 1985;21:181-3
- Yeh TE, Achanti B, Patel H, Pildes RS. Indomethacin therapy in premature infants with patent ductus arteriosus – determination of therapeutic plasma levels. *Dev Pharm Ther* 1989;12:169-78
- Weninger M, Pollak A, Salzer-Muhar U, Vergesslich KA, Salzer HR. Pharmacokinetics of intra-arterial indomethacin treatment for patent ductus arteriosus. *Eur J Pediatr* 1989;149:138-40
- Gal P, Ransom JL, Schall S, Weaver RL, Bird A, Brown Y. Indomethacin for patent ductus arteriosus closure. Application of serum concentrations and pharmacodynamics to improve response. *J Perinatol* 1990;10:20-6
- Robie DK, Waltrip T, Garcia-Prats JA, Pokorny WJ, Jaksic T. Is surgical ligation of a patent ductus arteriosus the preferred initial approach for the neonate with extremely low birth weight? *J Pediatr Surg* 1996;31:1134-7
- Trus T, Winthrop AL, Pipe S, Shah J, Langer JC, Lau GY. Optimal management of patent ductus arteriosus in the neonate weighing less than 800 g. *J Pediatr Surg* 1993;28:1137-9
- Palder SB, Schwartz MZ, Tyson KR, Marr CC. Management of patent ductus arteriosus: a comparison of operative v pharmacologic treatment. *J Pediatr Surg* 1987;22:1171-4

32. Zerella JT, Spies RJ, Deaver DCD, Dailry WJ, Haple DC, Trump DS. Indomethacin versus immediate ligation in the treatment of 82 newborns with patent ductus arteriosus. *J Pediatr Surg* 1983;18:835-41
33. Clyman RI. Recommendations for the postnatal use of indomethacin: an analysis of four separate treatment strategies [editorial]. *J Pediatr* 1996;128(5 Pt 1):601-7
34. Cotton RB, Stahlman MT, Bender HW, Graham TP, Catterton WZ, Kovar I. Randomized trial of early closure of symptomatic patent ductus arteriosus in small preterm infants. *J Pediatr* 1978;93:647-51
35. Fowlie PW. Prophylactic indomethacin: systematic review and meta-analysis. *Arch Dis Child, Fetal and Neonatal Edition* 1996;74:F81-F87
36. Ment LR, Oh W, Ehrenkranz RA, et al. Low-dose indomethacin and prevention of intraventricular hemorrhage: a multicenter randomized trial [see comments]. *Pediatrics* 1994;93:543-50
37. Hammerman C, Aramburo MJ. Prolonged indomethacin therapy for the prevention of recurrences of patent ductus arteriosus. *J Pediatr* 1990;117:771-6
38. Rennie JM, Cooke RW. Prolonged low dose indomethacin for persistent ductus arteriosus of prematurity [see comments]. *Arch Dis Child* 1991;66(1 Spec No):55-8
39. Rhodes PG, Ferguson MG, Reddy NS, Joransen JA, Gibson J. Effects of prolonged versus acute indomethacin therapy in very low birth-weight infants with patent ductus arteriosus. *Eur J Pediatr* 1988;147:481-4
40. Tammela O, Ojala R, Iivainen T, et al. Short versus prolonged indomethacin therapy for patent ductus arteriosus in preterm infants. *J Pediatr* 1999;134:552-7
41. Evans DH, Levene MI, Archer LN. The effect of indomethacin on cerebral blood-flow velocity in premature infants. *Dev Med Child Neurol* 1987;29:776-82
42. Lundell BP, Sonesson SE, Sundell H. Cerebral blood flow following indomethacin administration. *Dev Pharmacol Ther* 1989;13(2-4):139-44
43. Colditz P, Murphy D, Rolfe P, Wilkinson AR. Effect of infusion rate of indomethacin on cerebrovascular responses in preterm neonates. *Arch Dis Child* 1989;64(1 Spec No):8-12
44. Ohlsson A, Bottu J, Govan J, Ryan ML, Fong K, Myhr T. Effect of indomethacin on cerebral blood flow velocities in very low birth weight neonates with a patent ductus arteriosus. *Dev Pharmacol Ther* 1993;20(1-2):100-6
45. Hammerman C, Glaser J, Schimmel MS, Ferber B, Kaplan M, Eidelman AI. Continuous versus multiple rapid infusions of indomethacin: effects on cerebral blood flow velocity. *Pediatrics* 1995;95:244-8
46. Liem KD, Hopman JC, Kollee LA, Oeseburg B. Effects of repeated indomethacin administration on cerebral oxygenation and haemodynamics in preterm infants: combined near infrared spectrophotometry and Doppler ultrasound study [published erratum appears in *Eur J Pediatr* 1994 Oct;153:782]. *Eur J Pediatr* 1994;153:504-9
47. Mosca F, Bray M, Lattanzio M, Fumagalli M, Tosetto C. Comparative evaluation of the effects of indomethacin and ibuprofen on cerebral perfusion and oxygenation in preterm infants with patent ductus arteriosus. *J Pediatr* 1997;131:549-54
48. Patel J, Roberts I, Azzopardi D, Hamilton P, Edwards AD. Randomized double-blind controlled trial comparing the effects of ibuprofen with indomethacin on cerebral hemodynamics in preterm infants with patent ductus arteriosus [see comments]. *Pediatr Res* 2000;47:36-42
49. Bandstra ES, Montalvo BM, Goldberg RN, et al. Prophylactic indomethacin for prevention of intraventricular hemorrhage in premature infants. *Pediatrics* 1988;82:533-42
50. Bada HS, Green RS, Pourcyrous M, et al. Indomethacin reduces the risks of severe intraventricular hemorrhage [see comments]. *J Pediatr* 1989;115:631-7
51. Ment L, Vohr B, Allan W, et al. Outcome of children in the indomethacin intraventricular haemorrhage prevention trial. *Pediatrics* 2000;105:485-91
52. Kang NS, Yoo KH, Cheon H, et al. Indomethacin treatment decreases renal blood flow velocity in human neonates. *Biol Neonate* 1999;76:261-5
53. van Bel F, Guit GL, Schipper J, van de Bor M, Baan J. Indomethacin-induced changes in renal blood flow velocity waveform in premature infants investigated with color Doppler imaging. *J Pediatr* 1991;118(4 (Pt 1)):621-6
54. Seyberth HW, Rascher W, Hackenthal R, Wille L. Effect of prolonged indomethacin therapy on renal function and selected vasoactive hormones in very-low-birth-weight infants with symptomatic patent ductus arteriosus. *J Pediatr* 1983;103:979-84
55. Davis JM, Hendricks-Munoz KD, Hagberg D, Manning JA. The effects of indomethacin on renal function and intracranial hemorrhage in infants with patent ductus arteriosus. *Dev Pharmacol Ther* 1990;14:15-9
56. Yeh TF, Wilks A, Singh J, Betkerur M, Lilien L, Pildes RS. Furosemide prevents the renal side effects of indomethacin therapy in premature infants with patent ductus arteriosus. *J Pediatr* 1982;101:433-7
57. Brion LP, Campbell DE. Furosemide in indomethacin-treated infants – systematic review and meta-analysis. *Pediatr Nephrol* 1999;13:212-8
58. Baenziger O, Waldvogel K, Ghelfi D, Arbenz U, Fanconi S. Can dopamine prevent the renal side effects of indomethacin? A prospective randomized clinical study. *Klin Padiatr* 1999;211:438-41
59. Grosfeld JL, Chaet M, Molinari F, et al. Increased risk of necrotizing enterocolitis in premature infants with patent ductus arteriosus treated with indomethacin. *Ann Surg* 1996;224:350-5; discussion 355-7
60. Alpan G, Eyal F, Vinograd I, et al. Localized intestinal perforations after enteral administration of indomethacin in premature infants. *J Pediatr* 1985;106:277-81
61. Van Bel F, Van Zoeren D, Schipper J, Guit GL, Baan J. Effect of indomethacin on superior mesenteric artery blood flow velocity in preterm infants. *J Pediatr* 1990;116:965-70
62. Coombs RC, Morgan ME, Durbin GM, Booth IW, McNeish AS. Gut blood flow velocities in the newborn: effects of patent ductus arteriosus and parenteral indomethacin [see comments]. *Arch Dis Child* 1990;65:1067-71
63. Friedman Z, Whitman V, Maisels MJ, Berman W, Jr, Marks KH, Vesell ES. Indomethacin disposition and indomethacin-induced platelet dysfunction in premature infants. *J Clin Pharmacol* 1978;18(5-6):272-9

64. Ito K, Niida Y, Sato J, Owada E, Umetsu M. Pharmacokinetics of mefenamic acid in preterm infants with patent ductus arteriosus. *Acta Paediatr Jpn* 1994;36:387-91
65. Amato M, Huppi P, Markus D. Prevention of symptomatic patent ductus arteriosus with ethamsylate in babies treated with exogenous surfactant. *J Perinatol* 1993;13:2-7
66. Ng PC, So KW, Fok TF, Yam MC, Wong MY, Wong W. Comparing sulindac with indomethacin for closure of ductus arteriosus in preterm infants. *J Paed Child Health* 1997;33:324-8
67. Varvarigou A, Bardin CL, Beharry K, Chemtob S, Papageorgiou A, Aranda JV. Early ibuprofen administration to prevent patent ductus arteriosus in premature newborn infants. *JAMA* 1996;275:539-44
68. Van Overmeire B, Follens I, Hartmann S, Creten WL, Van Acker KJ. Treatment of patent ductus arteriosus with ibuprofen [see comments]. *Arch Dis Child, Fetal and Neonatal Edition* 1997;76:F179-F184
69. Pezzati M, Vangi V, Biagiotti R, Bertini G, Cianciulli D, Rubaltelli FF. Effects of indomethacin and ibuprofen on mesenteric and renal blood flow in preterm infants with patent ductus arteriosus. *J Pediatr* 1999;135:733-8
70. Van Overmeire B, Smets K, Lecoutere D, et al. A comparison of ibuprofen and indomethacin for closure of patent ductus arteriosus. *N Engl J Med* 2000;343:674-81