

Approaches to Studying the Development of Drug Metabolism in Children

Trevor N Johnson

Academic Unit of Molecular Pharmacology and Pharmacogenetics, Division of Clinical Sciences, University of Sheffield, L Floor, Royal Hallamshire Hospital, Sheffield S10 2JF, UK Tel +44 (0)114 2712984, Fax +44 (0)114 2720275, Email: t.n.johnson@sheffield.ac.uk

Altered drug disposition in the developing child occurs as a result of both biochemical and physiological changes. The clearance of many drugs is dependent on their biotransformation in the liver and small bowel and consequently is developmentally determined by a number of factors including both the activity and abundance of enzymes involved in Phase 1 & 2 drug metabolism. This paper reviews the *in vitro* and *in vivo* methods used to study the development of drug metabolism in humans with predominant reference to cytochrome P450 and UDP-glucuronyltransferase. The future possibility of *in silico* prediction of drug metabolism in children is also explored.

Paed Perinatal Drug Therapy 2002; 5: 75–92

Keywords: Drug metabolism – children – cytochrome P450 – glucuronidation – ontogeny

Introduction

From birth onwards changes in the pharmacokinetics and pharmacodynamics of administered drugs occur as a consequence of changes in body composition and maturation of organs and enzymes. Accordingly, effective and safe drug therapy in neonates, infants, children and adolescents requires a thorough understanding of human developmental biology and the ontogeny of the processes that govern the absorption, distribution, metabolism, excretion and action of drugs¹.

A number of factors contribute to the developmental changes in metabolic drug clearance in children. These include relative liver size, liver blood flow, extent of protein binding and the maturation of drug metabolising enzymes.

Altered drug metabolism can lead to the development of adverse effects in neonates and small infants that are not generally seen in the adult population. The altered metabolism of sodium valproate in children under 3 years of age is thought to be responsible for a higher

incidence of hepatotoxicity². The impaired metabolism of chloramphenicol in neonates can result in the grey baby syndrome (cyanosis and respiratory failure)³. Metabolic acidosis following the use of propofol in the critically ill child^{4,5} may be due to altered drug metabolism.

This review will focus principally on approaches and problems in determining the development of drug metabolism in humans with particular reference to Phase 1 reactions mediated by cytochromes P450 (CYP) and Phase 2 glucuronidation.

Enzymes involved in drug metabolism

Biotransformation generally converts lipophilic pharmacologically active drug molecules into polar metabolites that are then eliminated by the kidneys or other organs.

The liver is the major site of drug metabolism with intestinal drug extraction playing a secondary yet potentially important role in the pre-systemic elimination of orally administered drugs.

The process of drug metabolism is normally divided into two phases, phase 1 or functionalisation reactions and phase 2 conjugation reactions. Phase 1 reactions introduce or expose a chemically reactive group on which the phase 2 reactions can occur. The phase 2 reaction is usually the true detoxification pathway resulting in products that are generally water soluble and easily excreted. CYPs catalysing the mixed-function oxidation of xenobiotics comprise the most important group of enzymes involved in phase 1 metabolism. Others include the flavin-containing mono-oxygenases (FMO). Common phase 2 or conjugation reactions include glucuronidation, glycosidation, sulphation, methylation, acetylation and glutathione conjugation⁶.

Cytochrome P450

The CYP isoenzymes are a superfamily of haemoproteins that are the terminal oxidases of the mixed function oxidase system found in the endoplasmic reticulum. CYPs play two important roles in the living organism. Firstly, in the biosynthesis or biodegradation of endogenous compounds (steroids, fatty acids, prostaglandins) and, secondly in the oxidative biotransformation of exogenous compounds (drugs, environmental pollutants, toxins).

Approximately 500 CYP enzymes have so far been isolated spread across different species and grouped according to their amino acid sequences⁷. An amino acid homology greater than 40% defines a family, and greater than 55% defines a sub-family. The symbol CYP is used to denote both human and rat cytochrome P-450.

CYP is followed by an Arabic numeral denoting the family (CYP2), a capital letter designating the subfamily (CYP2C) and then another Arabic numeral representing the individual gene or enzyme (CYP2C9). The important human isoforms of the enzyme include CYPs 1A1 & 2, 2B6, 2C8–10, 2C19, 2D6, 2E1 and 3A4 & 5. The CYP3A subfamily are the most abundant enzymes in human liver and small intestine and are involved in the metabolism of around 50% of drugs. The principal substrates, inhibitors and inducers of the major human forms of this enzyme are summarised in Table 1.

Glucuronidation

Glucuronidation is an important detoxification pathway in humans. Many therapeutic drugs and their metabolites are substrates for UDP-glucuronyltransferases (UGT) leading to the formation of usually inactive glucuronides (a

notable exception is morphine 6 glucuronide which has analgesic activity⁹) which are then eliminated via the bile or urine¹⁰.

Individual UGT enzymes are defined according to the family (1 or 2 in humans), subfamily (A or B) and an arabic numeral representing the individual gene product. The liver represents the major site of glucuronidation in humans where five UGT1A and five UGT2B genes are expressed and define capacity^{11,12}. A number of UGT enzymes are expressed in extrahepatic sites^{12,13}. The important human hepatic enzymes are indicated in Table 2.

Approaches to studying the development of drug metabolism in humans

1. Human *in vivo* approaches

Probe substrates for specific CYP enzymes

A number of *in vivo* substrate probes for specific CYPs have been used and are summarised in Table 3.

One of the main problems with many of the tests relate to difficulties in their application for use in paediatric patients. Many of the tests involve the administration of radioactive substrates or intravenous drugs followed by the collection of multiple blood samples. This makes it more difficult to gain ethical approval for their use in children. Even when this is not the case such as with the caffeine breath test and urinary 6 β hydroxycortisol:cortisol ratio, there is usually a lack of validation of the tests for use in children. A summary of the consensus view as to the most appropriate *in vivo* probe for a specific CYP are shown in Table 3 with particular reference to those applicable to a paediatric population.

Caffeine breath test

The application of this test to assessing CYP1A2 activity in children has recently been reviewed³². Briefly, the caffeine breath test involves the oral administration of a non-radioactive stable isotope of caffeine (¹³C on the 3-methyl group). The exhaled labelled ¹³CO₂ correlates with CYP1A2 activity³³. The test is relatively non invasive and suitable for use in young children^{15,20}. A number of investigators have used this test in children to study the effects of disease states and drug interactions^{15–18}. Pons *et al.*¹⁹ have demonstrated no detectable changes in expired CO₂ from basal levels in neonates and young infants, whereas changes were measurable in all infants older than

Table 1. Substrates, inducers and inhibitors of each of the major human CYP isozymes⁸

Enzyme	Substrate	Inducer	Inhibitor
CYP1A1 (Extra-hepatic)	Chlorinated benzenes	Polycyclic hydrocarbons	Propofol
CYP1A2	Caffeine Erythromycin Haloperidol Paracetamol R-Warfarin Theophylline Tricyclic antidepressants	Hydrocarbons Omeprazole Phenobarbitone, Phenytoin, Polycyclic	Quinolone antibiotics
CYP2B6	Cyclophosphamide Mianserin Testosterone	None known	Orphenadrine
CYP2C8	Carbamazepine Diazepam Tricyclic antidepressants.	Phenobarbitone Rifampicin	Cimetidine (Not selective)
CYP2C9/10	Diclofenac Phenytoin S-Warfarin Tolbutamide	Rifampicin	Sulfaphenazole
CYP2C19	Diazepam Mephenytoin Tricyclic antidepressants	Phenobarbitone, Rifampicin	Fluconazole Omeprazole Warfarin
CYP2D6	Amitriptyline β -blockers Clozapine Codeine Debrisoquine SSRIs Tricyclic antidepressants	None Known	Amiodarone Flecainide Fluoxetine Quinidine Trifluoperidol
CYP2E1	Caffeine Chlorzoxazone Isoflurane Paracetamol	Benzene Ethanol Isoniazid	Dimethyl sulphoxide Disulfiram
CYP3A4	Carbamazepine Cisapride Clonazepam, Cyclosporin Diazepam Erythromycin Ethosuximide Etoposide Fentanyl Ketoconazole Lignocaine, Midazolam Nifedipine Ondansetron Rifampicin Ritonavir	Carbamazepine Glucocorticoids Omeprazole Phenobarbitone Phenytoin Rifampicin	Grapefruit juice Propofol Naringenin Triazole antifungals Troleandomycin
CYP3A5	Caffeine Cyclosporin Diltiazem Testosterone,	Dexamethasone	Troleandomycin
CYP3A7 (Fetal form)	Midazolam Testosterone		

Table 2. Endogenous and exogenous substrates of the major human UGT isozymes (de Wildt <i>et al.</i> ¹⁴ with modifications)			
UGT isoform	Endogenous substrate	Exogenous substrate	Comment
1A1*	Bilirubin	Ethinylloestradiol syndrome	↓ in Gilberts
1A3	Estrone	Norbuprenorphine	
1A4	Androstanediol	Amitriptyline Imipramine	
1A6*		1-Napthol 2-Napthol Naftazone Naproxen Paracetamol	
1A9	Estrone Phenol Propofol	Paracetamol	
1A10		Mycophenolic acid	
2B			Delayed onset associated with grey baby syndrome
2B4	Hyodeoxycholic		
2B7	Androsterone Epitestosterone	Buprenorphine Codeine Ibuprofen Morphine Naproxen Propranolol Valproic acid	
2B15*	Androgen steroids 4-Hydroxy-biphenyl	Eugenol	
2B17	Androsterone Dihydrotestosterone Testosterone		

* Polymorphic expression

33 days postnatal age and 45 weeks post-conceptual age, in agreement with *in vitro* findings. Lambert *et al.*²⁰ have demonstrated an increased 2 hour expired CO₂ in pre-pubertal children compared to adults which is contrary to the *in vitro* findings for the development of CYP1A2 in children and may reflect changes in relative liver size³⁴. There may be a number of potential problems with the caffeine breath test in the *in vivo* assessment of CYP1A2 activity. Children should be kept still during the test to avoid increased CO₂ production, which may have a dilutional effect on the test¹⁷.

The N3- and N7-demethylation of caffeine are the two mayor pathways of caffeine metabolism

in young infants and maturation of N1-demethylation occurs after 19 months³⁵. In the fetus the N3 and N7- reactions appear to be mediated by CYP3A rather than CYP1A2³⁶ with a switch over to CYP1A2 in the neonatal-infant stage. Full caffeine 3-demethylation is not observed until between 4 to 6 months of age³⁷. Thus, the changes in the pattern of metabolite formation and enzyme involvement in caffeine metabolism with age have to be considered when using this test for the determination of *in vivo* CYP1A2 development in children. Parker *et al.*¹⁸ suggest that the caffeine breath test is unsuitable for use in infants less than 6 months of age. The stable isotope is expensive.

Table 3. Recommended in vivo probe for CYPs (Tucker et al.³¹ with modifications)

CYP	Probe substrate(s)	Comments	Reference (paediatric studies)
1A2	Caffeine	Caffeine breath performed in children. Results only apply to a small number of drugs used in children	15–20
2B6	Bupropion	More validation required. No studies performed in children	
2C8	Unclear – Paclitaxel?	Cannot be given to healthy subjects.	
2C9	Tolbutamide	No studies performed in children using tolbutamide some evidence from placental transfer and prolonged half-life during first 2 days of postnatal life.	22
	Diclofenac		21
2C19	Omeprazole	Potential contamination from 3A4 pathway	23, 24
	Mephenytoin	Availability? No studies performed in children	
2D6	Debrisoquine	Availability and suitability for use in children?	
	Dextromethorphan	Alternative has been used in children but potential contamination from 3A4 pathway and urine pH-dependent renal excretion	25–27
2E1	Chlorzoxazone		
3A4	Midazolam (oral)	Not selective for 3A4 vs 3A5	28
	Midazolam (oral and iv)	Separates liver vs gut contributions; need for stable isotope labelling for concurrent oral and iv administration; staggered oral and iv dosing may avoid use of labelled drug	28
	Erythromycin	Erythromycin breath test marks 3A4 referentially to 3A5 in liver only. Results may be influenced by activity of P-glycoprotein transporter. Use of radioactive compound may be an issue especially in children	
	Midazolam (Oral) + Erythromycin (iv)	Suggested as a combined measure of both intestinal and hepatic CYP3A activity	
	6 β hydroxycortisol : cortisol	Has been used in children. Non invasive but results influenced by renal CYP3A5	29, 30

Diclofenac

Diclofenac is metabolised by CYP2C9 to 4-hydroxydiclofenac. Diclofenac is routinely used for analgesia in children. In a pharmacokinetic study of 500 mcg/kg diclofenac administered intravenously the plasma weight-corrected clearance was higher in children aged 4–6 years (7.7 ml/min/kg) compared to adults²¹ (3.3 ml/min/kg).

Omeprazole

Omeprazole is metabolised to 5-hydroxyomeprazole an index of CYP2C19 activity and to omeprazole sulphone an index of CYP3A4 activity

*in vivo*³⁸. Few studies have been undertaken in children. However, the clearance of intravenous omeprazole has been shown to be similar to adults in paediatric patients aged 4 months to 19 years²³. The oral clearance of omeprazole has been shown to be increased in children age 1–6 years compared to adults²⁴.

Mephenytoin

Mephenytoin has been used as a probe substrate to determine CYP2C19 phenotype in humans. (S)-(R)-mephenytoin undergoes rapid hydroxylation to form (S)-4 hydroxy mephenytoin (SOHP) with the slower formation of (R)- ethylhydantoin (PEH). The ratio of SOHP to PEH is used as a measure of CYP2C19 activity³⁹.

The test in adults involves the administration of 100mg mephenytoin followed by the collection of urine for 8 hours post dose^{40,41}. Although mephenytoin has been administered to children⁴² this test has so far not been applied to a paediatric population possibly because of worries over toxicity^{42,43}. Otherwise the test is relatively non-invasive. Stability issues have been raised regarding the assessment of CYP2C19 activity based on the SOHP : PEH ratio⁴⁴.

Dextromethorphan

Dextromethorphan is metabolised by CYP2D6 to dextrophan and by CYP3A to 3-methoxymorphinan^{45,46}. Dextromethorphan O-demethylation is well established as a measure of CYP2D6 *in vivo* and is highly correlated with CYP2D6 expression in human liver microsomes⁴⁷. The urinary ratio of dextromethorphan : dextrophan is most commonly used as an index of CYP2D6 activity and previous studies have shown a good correlation with other CYP2D6 probe substrates such as desipramine⁴⁸ and debrisoquine⁴⁹. In adults the test involves the administration of 25 to 30 mg of dextromethorphan and the collection of urine samples for 8 to 12 hours post dose^{41,50,51}.

The dextromethorphan test is relatively non-invasive, involves a relatively innocuous substrate and has been used to determine CYP2D6 polymorphism in children. Evans *et al.*²⁷ administered a dose of 30 mg to 26 children aged 3 to 21 years and collected urine for 4 hours post dose to determine debrisoquine oxidation phenotype.

The test has been applied to determining dextromethorphan phenotypes in paediatric patients with autoimmune hepatitis²⁶. In a recent study 300 mcg/kg dextromethorphan was administered to a group of neonates through the first year of life²⁵. Urine was collected into a nappy overnight. CYP2D6 phenotype consistent with genotype was achieved by 14.5 days postnatal age. The maturation of the CYP3A pathway (N-demethylation) was found to be delayed relative to CYP2D6. The use of this substrate in children is potentially problematic especially in neonates and infants because dextrophan formation as a measure of CYP2D6 is dependent on renal function⁵² which changes with developmental age. The ratio of dextromethorphan to dextrophan also depends on urinary pH.

Erythromycin breath test (ERBT)

The ERBT is one of the most extensively validated measures of *in vivo* CYP3A4 activity in the liver

in adults and was considered by some to be the gold standard^{53,54}. To date the test has not been applied to the paediatric population. The test is based on the CYP3A4 mediated N-demethylation of erythromycin producing formaldehyde which subsequently appears in the breath as CO₂⁵⁵. The test involves the intravenous administration of a small quantity of ¹⁴C labelled erythromycin, thereafter at timed intervals the subject blows into scintillation vials containing a CO₂ binding agent. The rate of production of radiolabel is then measured and expressed as the percentage of administered radiolabel exhaled during the first hour after injection⁵⁵.

The ERBT has been shown to predict the steady state trough blood levels of cyclosporin⁵⁶ and to predict its oral clearance⁵⁷. Lown *et al.*⁵⁴ showed the positive correlation of the ERBT with the weight adjusted clearance of midazolam (another *in vivo* probe of CYP3A) and McCrea *et al.*⁵⁸ have suggested the use of oral midazolam and the ERBT as a combined probe for hepatic and enterocytic CYP3A activity.

More recently doubt has been expressed as to whether the ERBT is a useful predictor of the clearance of some CYP3A drug substrates. In contrast to Lown *et al.*⁵⁴, Kinirons *et al.*⁵⁹ have failed to show correlation between midazolam clearance and the ERBT, likewise a lack of correlation between alfentanil clearance and the ERBT and has been observed by Krivoruk *et al.*⁶⁰. A possible reason for the discrepancy may be the fact that erythromycin⁶¹ but not midazolam or alfentanil⁶² is also a substrate for P-glycoprotein. This would result in lower intrahepatic concentrations of radiolabelled erythromycin and ultimately lower ¹⁴CO₂ exhalation⁶³. Another reason why the ERBT may not predict midazolam clearance is that erythromycin is a substrate for CYP3A4 but not CYP3A5⁶⁴ whereas midazolam is a substrate for both⁶⁵. Hence the ERBT will underestimate overall CYP3A activity in the 20 to 30% of patients with significant CYP3A5 hepatic levels⁶⁶.

Other disadvantages of the ERBT include: (i) the use of an intravenous radiolabelled substrate which will preclude its use especially in the paediatric population, (ii) the test does not measure intestinal CYP3A4 activity, (iii) only the rate of demethylation during the first hour is measured, interindividual differences in protein binding or volume of distribution could significantly affect the test result. Overall the ERBT may have limited application in the prediction of the CYP3A mediated metabolism of another substrate but may be useful for exploring the time course of induction or inhibition associated with

drug interactions⁶⁷. There has been some recent debate as to whether the CYP3A N-demethylation of erythromycin is the rate-limiting step in ¹⁴CO₂ production⁶⁸.

Midazolam

Midazolam is chiefly metabolised to 1' and 4'-hydroxymidazolam by CYP3A4 and CYP3A5⁶⁵. To perform the test an intravenous injection of 15 mcg/kg midazolam is administered and multiple blood samples collected up to 6 hours post dose. Drug and metabolites are measured by GC-MS or HPLC-MS.

Midazolam appears to have some advantages over the ERBT: it is metabolised by both CYP3A4 and CYP3A5 and is not a P-gp substrate. Furthermore administration of the drug intravenously and then orally may facilitate an overall assessment of intestinal and hepatic CYP3A⁶⁹. Further studies are required on the optimal use of midazolam as an oral CYP3A probe. The main disadvantages of the test are the administration of a clinically sedating dose of midazolam and the need for multiple blood samples. Rather than formally applying this test to children, studies built around the clinical use of midazolam have been applied to assessing the development of CYP3A activity.

The weight corrected plasma clearance of iv midazolam was significantly reduced in neonates compared to adults, 110–130 ml/h/kg and 380–660 ml/h/kg respectively^{70,71} and was even lower in preterm infants younger than 39 weeks gestation, 72–96 ml/h/kg^{70,72}. Because midazolam is only metabolised by CYP3A7 to a small extent⁶⁵, reduced clearance may be explained by low CYP3A4 activity following birth. The surge in CYP3A4 expression after birth is not mirrored by a large increase in midazolam clearance at this stage^{73,74}. Another study by Hughes *et al.*⁷⁵ showed lower weight corrected clearance in younger children up to 2 years of age (140–180 ml/h/kg) compared to children aged 3–13 years (780 ml/h/kg) who in turn had higher weight corrected clearance compared to adults (380 to 660 ml/h/kg). In a recent study both the i.v and oral clearance of midazolam were found to be increased (3,000 and 680 ml/h/kg respectively) in children age 6 months to <2 years compared to children age 2 to <12 (2,500 and 600 ml/h/kg respectively) and adolescents 12 to <16 (1500 and 560 L/h/kg respectively)²⁸.

Urinary 6β-hydroxycortisol

This is a simple non-invasive test that is easily performed even in a neonatal population. The ratio of 6β hydroxycortisol to free cortisol is

measured in a spot urine sample.

This ratio has been proposed as a measure of hepatic mixed function oxidase activity⁷⁶ and subsequently CYP3A activity⁷⁷. However, there is little evidence to support this. There is lack of correlation between the ERBT and 6βOHC:C ratio^{78,79}. Cortisol is also a substrate for P-gp which may in addition to the influence of renal CYP3A⁵⁸⁰ explain the lack of correlation between the 6βOHC:C urinary ratio and midazolam clearance. At best the 6βOHC:C ratio may be useful for looking at enzyme induction^{81–83}. A number of studies have applied this test to children. The 6βOHC:C ratio is higher in term compared to premature neonates³⁰. In infants 1 to 12 months of age the mean 6βOHC:C ratio was lower compared with that of neonates and adults²⁹.

Further studies are required on the use of probe substrates in determining *in vivo* CYP3A activity in man, especially with respect to their oral administration and with regard to the influence of P-gp. The tests require further validation for use in children.

Drugs in use

Phase 1

The pharmacokinetics of a number of drugs in routine clinical use have been applied to assessing the *in vivo* development of drug metabolism in children. This method is useful where the drug in question is predominantly metabolised by a single CYP enzyme. Ethical approval for such studies is generally easier to obtain compared to administering a probe substrate especially where iv access is already available and the number of blood samples to be collected is small. A number of approaches have been used to minimise the number of blood samples needed for such studies. These include sparse data pharmacokinetics, (small number of blood samples from a larger group of patients)^{84,85} and the use of steady state pharmacokinetics⁷⁵. A disadvantage with this approach is that it is opportunistic and thus more limited in its application.

A number of drugs routinely used in paediatric clinical practice and adopted as probes for specific CYP enzymes have already been mentioned including midazolam, diclofenac and omeprazole. Other drugs used in routine clinical practice that have been used to assess age-related changes in CYP activity include: for CYP3A: cyclosporin, tacrolimus, carbamazepine (also CYP2C8), etoposide, nifedipine, cisapride and for CYP2C9: phenytoin.

Phase 2

Morphine is largely metabolised by UGT2B7 to morphine-6-glucuronide and morphine-3-glucuronide⁹ and has been suggested as probe substrate for this UGT isoform⁸⁶. A number of *in vivo* studies have already been described on the developmental changes in morphine clearance in children⁸⁷⁻⁹⁰.

Paracetamol is mainly metabolised by UGT1A6 (and to a lesser extent by UGT1A9). The rate of paracetamol glucuronidation is negligible in the fetus⁹¹, low after birth and does not reach adult values before 10 years of age. Paracetamol is not the ideal probe for assessing UGT1A6 activity in children because of the compensatory higher sulphotransferase activity of the younger ages⁹².

Propofol has been proposed as an alternative probe for UGT1A9 activity⁸⁶. However, propofol is a high extraction drug with clearance being dependent mainly on blood flow rather than enzyme activity, thus, limiting its application as a probe substrate⁹³.

2. Human *in vitro* approaches

Several *in vitro* liver preparations are used in studying hepatic drug metabolism. These include liver slices, intact hepatocytes, S9 fractions and microsomes.

Liver slices and hepatocytes are useful for studying sequential oxidative and conjugative biotransformation in the same system as well as xenobiotic mediated induction⁹⁴. Many *in vitro* studies are done using human liver microsomes. Microsomes are closed vesicles of fragments of the endoplasmic reticular membrane and contain several drug metabolising enzymes including CYP, FMO and UDP-glucuronyl-transferase.

A number of investigators have used microsomes prepared from the livers of children at various developmental ages and then measured the expression and or the activity of specific CYP and other enzymes involved in drug metabolism.

The collection of liver biopsies is considered high risk and not justified for research purposes alone but may be acceptable when the primary purpose is for diagnosis or treatment⁹⁵. In practice, paediatric liver biopsy material is not only difficult to obtain for research purposes but when available is likely to have been collected from a patient with liver disease. Thus, many of the livers used to prepare microsomes have been obtained from aborted fetuses and post mortems in the first few hours after death^{96,97}. A recent study of the

developmental expression and activity of intestinal CYP3A4 was performed using duodenal and jejunal biopsies harvested at the same time as those to be used for clinical and diagnostic purposes⁹⁸. All human tissue used for *in vitro* studies should be characterised histologically for the presence of disease. Post mortem enzyme degradation may occur and tissue should be frozen at -80°C as soon as possible.

The expression of genes coding for specific CYP enzymes is often detected in fresh or frozen tissue by amplifying the gene in question using a reverse transcriptase polymerase chain reaction (RT-PCR) with appropriate primers.

Enzyme expression is usually measured by a technique such as Western blotting which requires separation of proteins by electrophoresis and the detection of the enzyme using specific antibodies.

Enzyme activity is determined by measuring the rate of metabolite appearance from a specific probe substrate. A number of model *in vitro* substrates have been defined for probing specific CYP enzyme activity (Table 4). Preferably more than one substrate should be used to investigate the ontogeny of CYP3A because of potential differences in the enzyme-substrate interaction^{99,100}.

To illustrate the application of *in vitro* techniques to investigate the ontogeny of specific CYP enzymes in the human liver, Sonnier and Cresteil³⁴ utilized liver tissue from stillborn or aborted fetuses (14–40 weeks), children aged 1 day to 9 years and from adult donors for kidney transplantation. Microsomes were prepared and CYP1A2 protein was measured by western blotting using a rat polyclonal CYP1A1 antibody. Enzyme activity was determined by co-incubating the microsomes with methoxyresorufin. CYP1A2 expression and activity was absent in microsomes prepared from fetal and neonatal livers and increased in infants aged 1–3 months to attain 50% of the adult value at one year.

Problems with the human *in vitro* approach include the availability of suitable tissue samples, ethical constraints, especially since the Redfern report¹⁰¹, (The Report of The Royal Liverpool Children's Inquiry) the availability of isoform specific antibodies to assess expression by immunoblotting, size of the tissue samples available, the non-physiological nature of the preparations used and the availability of sufficiently sensitive assay systems to measure metabolite formation. The main advantage of this approach is that changes in true human enzymology with age can be ascertained.

Table 4. Recommended in vitro probe substrates ³¹		
CYP	Substrates	
	Preferred	Acceptable
1A2	Ethoxyresorufin Phenacetic	Caffeine Theophylline
2A6	Coumarin	
2B6	S-Mephenytoin	Bupropion (availability of metabolite standards?)
2C8	Paclitaxel	
2C9	Diclofenac S-Warfarin	Tolbutamide (low turnover)
2C19	Omeprazole S-Mephenytoin	
2D6	Bufuralol Dextromethorphan Metoprolol	Codeine Debrisoquine
2E1	Chlorzoxazone	4-nitrophenol Lauric acid
3A4	Midazolam Testosterone	Cyclosporin Erythromycin *

*Strongly recommend using at least two structurally unrelated substrates.

Developmental factors may influence the suitability of an *in vitro* probe substrate. For instance, caffeine is metabolised by CYP3A in fetal liver³⁶, whereas in adults it is predominantly metabolised by CYP1A2¹⁰².

Studies on the developmental expression of polymorphic enzymes such as CYP2D6 and CYP2C19 need to account for the numbers of poor metabolisers within each age group. This is itself potentially difficult as there is little information about the appearance of the different phenotypes during development¹⁰³. Peng *et al.*¹⁰⁴ studied the *in vitro* acetylation of 7-amino-clonazepam in human fetal and adult liver preparations. The adults but not the fetuses could be classified into fast and slow acetylators.

In general, enzyme expression and activity correlate closely in adult humans¹⁰⁵. However, during development this is not always the case. Strassberg *et al.*¹⁰⁶ have demonstrated an early phase in UGT development with the appearance of gene transcripts by 6 months followed by a later phase characterised by up regulation of UGT protein expression to levels found in adults by 2 years. Even by 2 years the activity of many UGT enzymes was significantly lower than in adults. The different stages of ontogeny are characterised by dynamic changes in gene expression. Thus, *in vitro* studies drawing conclusions based on a small

number of tissue samples, representing a narrow time window must be viewed with caution¹⁰⁷.

Investigations utilising microsomal and other subcellular liver and intestinal fractions in the study of the development of the major enzymes involved in drug metabolism are shown in Tables 5 and 6. Two reviews are available describing the *in vitro* ontogeny of human phase 1¹⁰⁷ and phase 2¹⁰⁸ drug metabolising enzymes.

3. Animal models

The relative ease of obtaining animals, usually rats, at the relevant stage of development and then harvesting the organs of interest, make the use of animal models attractive for studying the ontogeny of drug metabolism systems.

The development of rat hepatic CYP3A has been shown to be both age and sex dependent¹³⁰⁻¹³¹. Enzyme expression and activity increases up to day 25 and then continues to rise in male but falls away in female animals. The expression and inducibility of P450 enzymes during liver ontogeny has been reviewed by Rich and Boobis¹³². In contrast to rat hepatic CYP3A, small bowel CYP3A expression and activity is virtually absent until weaning (day 20) and then surges, reaching a plateau by 40 days sustained into adulthood¹³¹.

Table 5. In vitro study of developing phase 1 enzymes in humans								
Tissue and source	In vitro preparation	Enzyme CYP	Probe Substrate	Other methods	Relative % expression / activity			Reference
					Adult	Child (age)	Fetus	
L, AF, SS	Microsomes	1A		Western blotting	Detected (+++)		Not detected	109
AF, SS	Microsomes	1A1	Ethoxyresorufin -o-demethylase		0		100 (7-9wk)	110
L, AF, OD, B	MRNA and cDNA extract	1A1		RT-PCR	Detected (++)		Not detected	111
L, AF, SS,OD	Microsomes	1A2	Methoxyresorufin demethylation	Western blotting	100	50 (1 yr)	0	34
L, AF, OD, B	MRNA and cDNA extract	1A2		RT-PCR	Detected (+++)		Not detected	111
L, AF, PM, SS	Microsomes / RNA/section	1A2		Immunostaining, Western blotting, Northern blotting	100	53 (5yr)	Not detected	112
L, SS, B	Microsomes	2A6		Western blotting	100	<100 (1yr)		113
L, SS, B	Microsomes	2B6		Western blotting	100	10 (1yr)	-	113
L, AF, OD, B	MRNA and cDNA extract	2B6		RT-PCR	Detected (+++)		Not detected	111
L, AF, PM, SS	Microsomes / RNA/section	2C		Immunostaining, Western blotting, Northern blotting	100	47 (5d) 91 (5yr)	< 10	112
L, OD, AF, PM	Microsomes	2C 2C8 2C9 2C10 2C18	Tolbutamide	RT-PCR	100	5 (1 d) 20 (1 mo) 65 (7d)	1-2 50 0** 40	114, 115
					100			
					100	6 (7d) 25 (1mo) 85 (1d) 120 (1yr)		
					100			
					100			
L, AF, SS	Microsomes	2C		Western blotting	Detected (+++)		Detected (+)	109
L, AF, OD, B	MRNA and cDNA extract	2C8-19*		RT-PCR	Detected (++++)		Detected (++)	111

Tissue and source	In vitro preparation	Enzyme CYP	Probe Substrate	Other methods	Relative % expression / activity			Reference
					Adult	Child (age)	Fetus	
L, AF, OD, B	MRNA and cDNA extract.	2D6*		RT-PCR	Detected (+++)		Detected (+)	111
L, AF, OD	Microsomes	2D6*	Dextromethorphan		100	Detected RNA(++)	0	116
L, AF, OD	Microsomes	2D6*	Dextromethorphan	Western blotting, RNA analysis	100	15 (7 d) 50 (1 m)	2-5	117
L, AF, OD	Microsomes	2D6*	Dextromethorphan		100		< 5	118
L, AF, OD, B	MRNA and cDNA extract	2E1		RT-PCR	Detected (+++)		Not detected	111
L, AF, OD, SS	Microsomes	2E1	Ethanol oxidation	Western blotting, RT-PCR	100		12-27	119
L, AF, PM	Microsomes	2E1	Chlorzoxazone hydroxylation	Northern blotting	100	15 (1 mo) 80 (1 yr)	0	120
L, AF		2E1		RT-PCR	100		10-30	121
L, AF, OD, SS and olfactory tissue	Microsomes	2J2		Western blotting	100		Near 100	122
L, AF, PM	Microsomes	3A4/5*	DHEA hydroxylation, Testosterone hydroxylation	Western blotting	100	50 (6 mo) 120 (1y)	<10	96
L, AF, OD, B	MRNA and cDNA extract	3A4/5*		RT-PCR	Detected (+++)		Detected (++)	111
L, AF, PM	Microsomes MRNA and cDNA extract	3A7	DHEA	RT-PCR	<10 Detected (++)	150 (1d) 50 (8d)	100 Detected (++++)	96
Duodenum / Jejunum, B, AE, NS	Microsomes	3A4	Testosterone hydroxylation Dextromethorphan	Western blotting	100	50 (1mo) 75 (2 yr)	5	98
L, AF, PM, SS	Microsomes	3A4	Cisapride		56	100	3.5	123

* Polymorphism

** Approximately 5% detection of CYP2C9 RNA in fetus compared to adult

Tissue sources: AF, Aborted fetus; L, Liver, SS, Surgical section; OD, Organ donor; PM, Post mortem; B, Biopsy. DHEA, Dehydroepiandrosterone.

Table 6. <i>In vitro</i> studies with hepatic microsomes on the development of phase 2 glucuronidation enzymes in humans						
Source	Enzyme UGT	Probe substrate/ Other Method	Relative % expression / activity			Reference
			Adult	Child (age)	Fetus	
L, AF, OD	1A1*	Bilirubin	100	1 (1mo)	0.1	124
L, AF, SS, PM	1A1*	Bilirubin	100		<14	125
L, AF, PM	1A1*	Bilirubin/ Western blotting	100	<14 (term)	<14	126
L, AF, SS, PM	1A3		100		30	14
L, SS	1A4	Amitriptyline	100	5–10 (18mo)	<5	106
L, AF, SS, PM	1A6*	1-naphthol	100		<14	125
L, AF,OD	1A6*	2-naphthol	100		1	127
L, AF, SS, PM	1A9	Phenol	100		<14	125
L, AF, SS, PM	1A9	4-butylphenol	100		5	106
L, AF, SS, PM	2B7*	Androsterone	100		<14	125
L, AF, OD	2B7*	Morphine	100		10–20	128
L, OD,SS	2B7*	Buprenorphine	100		10	106
L, AF, SS, PM	2B17	Testosterone	100	<14		125
L, AF	2B17	Western blotting	Detected (+++)	Detected (+)		129

* Polymorphism

Tissue sources: AF, Aborted fetus; L, Liver; SS, Surgical section; OD, Organ donor; PM, Post mortem.

Although these data are of interest to the toxicologists there are a number of inherent problems in applying it to humans. There are differences in the expressed enzymes. In man the predominant CYP3A enzymes are CYP3A7 (fetal form), CYP3A4 and CYP3A5, whereas in rats they are CYP3A1, CYP3A2, CYP3A9 and CYP3A18¹³³. Although they are all involved in the metabolism of certain substrates e.g. testosterone, major differences exist. For instance, mephenytoin is metabolised by CYP3A in the rat but by CYP2C19 in humans^{134,135}. Many enzymes in the rodent have a sex specific ontogeny (CYP2C11, CYP2C12, CYP2C13, CYP3A2) whereas this is not the case in humans. Animal models also often ignore wide genetic polymorphisms which occur in humans such as CYP2D6 and CYP2C29¹³⁶.

Coupled with the contrasts in enzymology, there are often cross species developmental mismatches occurring on a more physiological level. This is especially so when investigating the development of drug metabolism in the small bowel. In rodents the bowel continues to develop after birth and only becomes functional by the

time of weaning¹³⁷. In contrast, the function of human bowel is ahead of need at birth¹³⁸. Neonates have around 50% of the small intestinal CYP3A expression and activity compared to adults⁹⁸. This contrasts with the newborn rat where the enzyme is absent.

Allometric scaling (AS) using *in vivo* data obtained in animal species has been used to attempt to predict pharmacokinetic (PK) parameters in humans¹³⁹. AS assumes that many physiological processes are a function of the size of the animal and, thus, PK parameters should be a function of total body weight¹⁴⁰. AS is therefore applied in a retrospective manner to compare the PK of a drug across several species and predict the parameters in humans¹³⁶. There are many problems in using AS to predict PK parameters in adult humans due to its empirical nature¹⁴¹, including interspecies differences in metabolism. Applying the technique to studying the metabolism of drugs in the developing human from PK data derived from developing animals would introduce yet more confounding factors and further reduce the possibility of useful predictions.

Overall, although animal models are of some relevance to acute toxicity testing of drugs during development, they are generally unreliable at predicting the ontogeny of drug metabolism in humans.

4. *In silico* approaches

Computer simulation is being employed increasingly within the pharmaceutical industry. The prediction of absorption, distribution, metabolism and excretion parameters for a specific drug are becoming possible from its chemical structure alone^{142,143}. An area of rapid development is the prediction of *in vivo* pharmacokinetics parameters from *in vitro* data. Such *in vitro-in vivo* extrapolation (IVIVE) depends on the availability of good quality *in vitro* enzyme kinetic data such as maximum enzyme velocity (V_{max}), Michaelis-Menten constant (K_m) for the drug substrate and inhibitor constant (K_i) and inhibitor concentration (I) for the inhibitor.

Computer simulation using SIMCYP a programme developed at the University of Sheffield and incorporating population variability has been successfully applied in the prediction of the variability of a number of drug-drug interactions including methadone-ritonavir¹⁴⁴, ketoconazole-midazolam¹⁴⁵, fluconazole-midazolam¹⁴⁶ and sildenafil-ritonavir¹⁴⁷.

One future development will be the *in silico* prediction of the likely pharmacokinetic parameters of drugs in children. The *in vitro* developmental patterns of many of the CYP enzymes in children are documented (see earlier) as are other key physiological parameters such as liver size, renal function and intestinal development. The first stage will be to simulate the changes and variability in clearance of a drug such as midazolam with age, and compare the results with those available in the literature. Additional *in vitro* data will be required to build a truly paediatric version of SIMCYP but once validated it will form a valuable tool both for the pharmaceutical industry and in the clinical setting.

Conclusions

An array of approaches exist for studying the development of drug metabolism in children each with their own advantages and disadvantages. Future studies are required not only to extend the current data on the development of specific enzyme systems with age but also to further validate methods for use in children. Drug metabolism is an important determinant of both drug dose and frequency of administration. Thus, it is important that information on the changes

in drug metabolism with age is accounted for in the clinical use of medicines for children. To this end, the development of validated *in silico* approaches may facilitate the reliable prediction of pharmacokinetic parameters for drugs to be used in children. This in turn may lead to a more scientific approach to dosage recommendation and the design of clinical studies in children.

An area that is currently generating research interest is the relationship between the ontogeny of drug metabolising enzymes and the pathogenesis of adverse drug reactions (ADRs) in children. Delayed maturation of enzymes systems over the first 2–4 weeks of life may contribute to the development of type A (dose dependent) ADRs. Conversely it is postulated that the increased metabolism and thus circulating metabolites of possibly reactive metabolites in young children may predispose to certain type B (Idiosyncratic, non-dose dependent) ADRs. Future research in this area must not only focus on the ontogeny of the drug metabolising enzymes and transporter proteins, but also other systems involved in the generation of ADRs such as the development of the immune system. Only by understanding the underlying mechanisms of ADRs in children can we ultimately begin to prevent them from occurring.

References

1. Morselli PL, Franco-Morselli R, Bossi L. Clinical pharmacokinetics in newborns and infants. Age-related differences and therapeutic implications. *Clin Pharmacokinet* 1980; 5: 485–527.
2. Fisher E, Siemes H, Pund R, Wittfoht W, Nau H. Valproate metabolites in serum and urine during antiepileptic therapy in children with infantile spasms; abnormal metabolite pattern associated with reversible hepatotoxicity. *Epilepsia* 1992; 33:165–171.
3. Weiss CF, Glasko AJ, Weston JK. Chloramphenicol in the newborn infant; *N Eng J Med* 1960; 262: 787–794.
4. Parke TJ, Stevens JE, Rice AS *et al.* Metabolic acidosis and fatal myocardial failure after propofol infusion in children: five case reports. *BM J* 1992; 305: 613–616.
5. Strickland RA, Murray MJ. Fatal metabolic acidosis in a pediatric patient receiving an infusion of propofol in the intensive care unit: is there a relationship? *Crit Care Med* 1995; 23: 405–9.
6. Gibson GG, Skett P. Introduction to drug metabolism, 2nd edition, Blackie Academic and Professional, London, 1997.
7. Nelson DR, Koymans L, Kamataki T *et al.* P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics* 1996; 6: 1–42.
8. Chang GMW, Kam PCA. The physiological and pharmacological roles of cytochrome P450 isoenzymes. *Anaesthesia* 1999; 54: 42–50.
9. Coffman BL, Rios GR, King CD *et al.* Human UGT2B7 catalyses morphine glucuronidation. *Drug Metab Dispos* 1997; 25: 1–4.

10. Dutton GJ, Glucuronidation of drugs and other compounds. Boca Raton, Florida: CRC Boca Raton Press, 1980.
11. Tukey RH, Strassburg CP. Human UDP-glucuronosyltransferases: metabolism, expression, and disease. *Ann Rev Pharmacol Toxicol* 2000; 40: 581–616.
12. Strassburg CP, Nguyen N, Manns MP *et al.* UDP-glucuronosyltransferase activity in human liver and colon. *Gastroenterology* 1999; 116: 149–160.
13. Strassburg CP, Kneip S, Topp J. Polymorphic gene expression and interindividual variation of UDP-glucuronosyltransferase activity in human small intestine. *J Biol Chem* 2000; 275: 36164–36171.
14. de Wildt SN, Kearns GL, Leeder JS, van den Anker JN. Glucuronidation in humans; Pharmacogenetic and developmental aspects. *Clinical Pharmacokinetics* 1999; 36: 439–452.
15. Parker AC, Preston T, Heaf D, Kitteringham NR, Choonara I. Inhibition of caffeine metabolism by ciprofloxacin in children with cystic fibrosis as measured by the caffeine breath test. *Br J Clin Pharmacol* 1994; 38: 573–576.
16. Parker AC, Pritchard P, Preston T, Dalzell AM, Choonara I. Lack of inhibitory effect of cimetidine on caffeine metabolism in children using the caffeine breath test. *Br J Clin Pharmacol* 1997; 43: 467–470.
17. Parker AC, Pritchard P, Preston T, Smyth RL, Choonara I. Enhanced drug metabolism in young children with cystic fibrosis. *Arch Dis Child* 1997; 77: 239–41.
18. Parker AC, Pritchard P, Preston T, Choonara I. Induction of CYP1A2 activity by carbamazepine in children using the caffeine breath test. *Br J Clin Pharmacol* 1998; 45: 176–178.
19. Pons G, Blais JC, Rey E *et al.* Maturation of caffeine N-demethylation in infancy: a study using the 13CO₂ breath test. *Pediatric Res* 1988; 23: 632–636.
20. Lambert GH, Schoeller DA, Kotake AN, Flores C, Hay D. The effect of age, gender, and sexual maturation on the caffeine breath test. *Dev Pharmacol Ther* 1986; 9: 375–88.
21. Korpela R, Olkkola KT. Pharmacokinetics of intravenous diclofenac sodium in children. *Eur J Clin Pharmacol* 1990; 38: 293–295.
22. Christesen HB, Melander A. Prolonged elimination of tolbutamide in a premature newborn with hyperinsulinaemic hypoglycaemia. *Eur J Endocrinol* 1998; 138: 698–701.
23. Jacqz-Aigrain E, Bellaich M, Faure C, Andre J, Rohrlisch P, Baudouin V, Navarro J. Pharmacokinetics of intravenous omeprazole in children. *Eur J Clin Pharmacol* 1994; 47: 181–185.
24. Andersson T, Hassall E, Lundborg P *et al.* Pharmacokinetics of orally administered omeprazole in children. International pediatric omeprazole pharmacokinetic group. *Am J Gastroenterol* 2000; 95: 3101–3106.
25. Leeder JS, Gotschall RR, Gaedigk A, Adcock K, Wilson JT, Kearns GL. Acquisition of functional CYP2D6 and CYP3A4 activities in vivo in the first year of life. *Drug Metab Rev.* 2000; 32(Suppl. 2): 281(abstrac #289).
26. Jacqz-Aigrain E, Laurent J, Alvarez F. Dextromethorphan phenotypes in paediatric patients with autoimmune hepatitis. *Br J Clin Pharmacol* 1990; 30:153–154.
27. Evans WE, Relling MV, Petros WP, Meyer WH, Mirro J, Crom WR. Dextromethorphan and caffeine as probes for simultaneous determination of debrisoquin-oxidation and N-acetylation phenotypes in children. *Clin Pharmacol Ther* 1989; 45: 568–573.
28. Reed MD, Rodarte A, Blumer JL, Khoo KC, Akbari B, Pou S, Kearns GL. The single dose pharmacokinetics of midazolam and its primary metabolite in pediatric patients after oral and intravenous administration. *J Clin Pharmacol* 2001; 41: 1359–1369.
29. Nakamura H, Hirai M, Ohmori S *et al.* Changes in urinary 6 β -hydroxycortisol/cortisol ratio after birth in human neonates. *Eur J Clin Pharmacol* 1998; 53: 343–346.
30. Vauzelle-Kervroedan F, Rey E, Pariente-Khayat A *et al.* Non-invasive in vivo study of the maturation of CYP 3A in neonates and infants. *Eur J Clin Pharmacol* 1996; 51: 69–72.
31. Tucker GT, Houston JB, Huang SM. Optimizing drug development: strategies to assess drug metabolism/transporter interaction potential-toward a consensus. *Clin Pharmacol Ther* 2001; 70: 103–14.
32. Webster E, McIntyre J, Choonara I, Preston T. The caffeine breath test and CYP1A2 activity in children. *Paed Perinatal Drug Ther* 2002; 5: 28–33.
33. Wietholtz , Voegelin M, Arnaud MJ, Bircher J, Preisig R. Assessment of the cytochrome P-448 dependent liver enzyme system by a caffeine breath test. *Eur J Clin Pharmacol* 1981; 21: 53–59.
34. Sonnier M, Cresteil T. Delayed ontogenesis of CYP1A2 in the human liver; *Eur J Biochem* 1998; 251: 893–898.
35. Carrier O, Pons G, Rey E *et al.* Maturation of caffeine metabolic pathways in infancy. *Clin Pharmacol Ther* 1988; 44:145–151.
36. Cazeneuve C, Pons G, Rey E. Biotransformation of caffeine in human liver microsomes from fetuses, neonates, infants and adults. *Br J Clin Pharmacol* 1994; 37: 405–412.
37. Aranda JV, Collinge JM, Zinman R, Watters G. Maturation of caffeine elimination in infancy. *Arch Dis Childhood* 1979; 54: 946–949.
38. Tassaneeyakul W, Vannaprasaht S, Yamazoe Y. Formation of omeprazole sulphone but not 5-hydroxyomeprazole is inhibited by grapefruit juice. *Br J Clin Pharmacol* 2000; 49: 139–144.
39. Wilkinson GR, Guengerich FP, Branch RA. Genetic polymorphism of S-mephenytoin hydroxylation. *Pharmacol Ther.* 1989; 43: 53-76.
40. Eap CB, Bondolfi G, Zullino D *et al.* Pharmacokinetic drug interaction potential of risperidone with cytochrome p450 isozymes as assessed by the dextromethorphan, the caffeine, and the mephenytoin test. *Ther Drug Monit* 2001; 23: 228–231.
41. Eap CB, Guentert TW, Schaublin-Loidl M *et al.* Plasma levels of the enantiomers of thioridazine, thioridazine 2-sulfoxide, thioridazine 2-sulfone, and thioridazine 5-sulfoxide in poor and extensive metabolizers of dextromethorphan and mephenytoin. *Clin Pharmacol Ther* 1996; 59: 322–331.
42. Ilyes I, Tornai A, Kirilina S, Gyorgy I. Influence of anticonvulsant drugs on thyroid hormones in epileptic children. *Acta Paediatrica Hungarica.* 1985; 26: 307–310.
43. Troupin AS, Ojemann LM, Dodrill CB. Mephenytoin: a reappraisal. *Epilepsia.* 1976; 17: 403–414.
44. Zhang Y, Blouin RA, McNamara PJ, Steinmetz J, Wedlund PJ. Limitation to the use of the urinary S-/R-mephenytoin ratio in pharmacogenetic studies. *Br J Clin Pharmacol* 1991; 31: 350–352.
45. Barnhart JW. The urinary excretion of dextromethorphan and three metabolites in dogs and humans. *Toxicol Appl Pharmacol* 1980; 55: 43–48.

46. Kerry NL, Somogli AA, Bochner F, Mikus G. The role of CYP2D6 in primary and secondary oxidative metabolism of dextromethorphan: in vitro studies using human liver microsomes. *Br J Clin Pharmacol* 1994; 38: 243–248.
47. Rodrigues AD, Kukulka MJ, Surber BW *et al.* Measurement of liver microsomal cytochrome p450 (CYP2D6) activity using [O-methyl-14C]dextromethorphan. *Anal Biochem* 1994; 219: 309–20.
48. Spina E, Gitto C, Avenoso A, Campo GM, Caputi AP, Perucca E. Relationship between plasma desipramine levels, CYP2D6 phenotype and clinical response to desipramine: a prospective study. *Eur J Clin Pharmacol* 1997; 51: 395–398.
49. Kupfer A, Schmid B, Preisig R, Pfaff G. Dextromethorphan as a safe probe for debrisoquine hydroxylation polymorphism. *Lancet* 1984; 2: 517–518.
50. Jones DR, Gorski JC, Haehner BD, O'Mara EM Jr, Hall SD. Determination of cytochrome P450 3A4/5 activity in vivo with dextromethorphan N-demethylation. *Clin Pharmacol Ther* 1996; 60:374–784.
51. Streetman DS, Bleakley JF, Kim JS *et al.* Combined phenotypic assessment of CYP1A2, CYP2C19, CYP2D6, CYP3A, N-acetyltransferase-2, and xanthine oxidase with the "Cooperstown cocktail". *Clin Pharmacol Ther* 2000; 68: 375–383.
52. Rostami-Hodjegan A, Kroemer HK, Tucker GT. In-vivo indices of enzyme activity: the effects of renal impairment on the assessment of CYP2D6 activity. *Pharmacogenetics* 1999; 9: 277–286.
53. Kinirons MT, O'Shea D, Downing TE *et al.* Absence of correlations among three putative in vivo probes of human cytochrome P4503A activity in young healthy men. *Clin Pharmacol Ther* 1993; 54: 621–629.
54. Lown KS, Thummel KE, Benedict PE *et al.* The erythromycin breath test predicts the clearance of midazolam. *Clin Pharmacol Ther* 1995; 57: 16–24.
55. Watkins PB, Murray SA, Winkelman LG, Heuman DM, Wrighton SA, Guzelian PS. Erythromycin breath test as an assay of glucocorticoid-inducible liver cytochromes P-450. *J Clin Invest* 1989; 83: 688–697.
56. Watkins PB, Hamilton TA, Annesley TM, Ellis CN, Kolars JC, Voorhees JJ. The erythromycin breath test as a predictor of cyclosporin A blood levels. *Clin Pharmacol Ther* 1990; 48: 120–129.
57. Turgeon DK, Normolle DP, Leichtman AB, Annesley TM, Smith DE, Watkins PB. Erythromycin breath test predicts oral clearance of cyclosporin A in renal transplant recipients. *Clin Pharmacol Ther* 1992; 52: 471–478.
58. McCrea J, Prueksaritanont T, Gertz BJ. Concurrent administration of the erythromycin breath test (EBT) and oral midazolam as in vivo probes for CYP3A activity. *J Clin Pharmacol* 1999; 39: 1212–1220.
59. Kinirons MT, O'Shea D, Kim RB *et al.* Failure of the erythromycin breath test to correlate with midazolam clearance as a probe of cytochrome P4503A. *Clin Pharmacol Ther* 1999; 66: 224–231.
60. Krivoruk Y, Kinirons MT, Wood AJJ, Wood M. Metabolism of cytochrome P4503A substrates in vivo administered by the same route: lack of correlation between alfentanil clearance and erythromycin breath test. *Clin Pharmacol Ther* 1994; 56: 608–614.
61. Schuetz EG, Yasuda K, Arimori K, Schuetz JD. Human MDR1 and mouse mdr1a P-glycoprotein after the cellular retention and disposition of erythromycin, but not of retinoic acid or benzo(a)pyrene. *Arch Biochem Biophys* 1998; 350: 340–347.
62. Kim RB, Fromm MF, Wandel C *et al.* The drug transporter P-glycoprotein limits oral absorption and brain entry of HIV-1 protease inhibitors. *J Clin Invest* 1998; 101: 289–294.
63. Kinirons MT, O'Shea D, Kim RB, Wood AJJ, Wilkinson GR. Erythromycin breath test (Letter reply). *Clin Pharmacol Ther* 2000; 67: 577–578.
64. Wrighton SA, Brian WR, Sari MA *et al.* Studies on the expression and metabolic capabilities of human liver cytochrome P-450IIIa5 (HLp3). *Mol Pharmacol* 1990; 38: 207–213.
65. Gorski JC, Hall SD, Jones DR, VandenBranden M, Wrighton SA. Regioselective biotransformation of midazolam by members of the human cytochrome P450 3A (CYP3A) subfamily. *Biochem Pharmacol* 1994; 47: 1643–1653.
66. Wrighton SA, Ring BJ, Watkins PB, Watkins PB, Vanderbranden M. Identification of a polymorphically expressed member of the human cytochrome P450 III family. *Mol Pharmacol* 1989; 36: 97–105.
67. Cheng CL, Smith E, Carver PL *et al.* Steady state pharmacokinetics of delavirdine in HIV-positive patients: effects on erythromycin breath test. *Clin Pharmacol Ther* 1997; 61: 531–543.
68. Chiou WL, Jeong HY, Wu TC, Ma C. Use of the erythromycin breath test for in vivo assessment of cytochrome P450 3A activity and dosage individualisation. *Clin Pharmacol Ther* 2001; 70: 305–339.
69. Thummel KE, O'Shea D, Paine MF *et al.* Oral first pass elimination of midazolam involves both gastrointestinal and hepatic CYP3A-mediated metabolism. *Clin Pharmacol Ther* 1996; 59: 491–502.
70. Burtin P, Jacqz-Aigrain E, Girard P. Population pharmacokinetics of midazolam in neonates. *Clin Pharmacol Ther* 1994; 56: 615–625.
71. Reves JG, Fragen RJ, Vinik HR, Greenblatt DJ. Midazolam pharmacology and uses. *Anesthesiology* 1985; 62: 310–324.
72. Harte GJ, Gray PH, Casteel HB, Steer PA, Charles BG. Haemodynamic responses and population pharmacokinetics of midazolam following administration to ventilated premature neonates. *J Paediatr Child Health* 1997; 33: 335–338.
73. Wells TG, Ellis EN, Casteel HB. Pharmacokinetics of a single dose midazolam in children. *Clin Pharmacol Ther* 1991; 49: 160.
74. Jacqz-Aigrain E, Daoud P, Burtin P, Maherzi S, Beaufils F. Pharmacokinetics of midazolam during continuous infusion in critically ill neonates. *Eur J Clin Pharmacol* 1992; 42: 329–332.
75. Hughes J, Gill AM, Mulhearn H, Powell E, Choonara I. Steady state plasma concentrations of midazolam in critically ill infants and children. *Ann Pharmacother* 1996; 30: 27–30.
76. Park BK. Assessment of urinary 6 beta-hydroxycortisol as an in vivo index of mixed function oxidase activity. *Br J Clin Pharmacol* 1981; 12: 97–102.
77. Waxman DJ, Attisano C, Guengerich FP, Lapenson DP. Human liver microsomal steroid metabolism: identification of the major microsomal steroid hormone 6 beta-hydroxylase cytochrome P-450 enzyme. *Arch Biochem Biophys* 1988; 263: 424–436.
78. Watkins PB, Turgeon DK, Saenger P *et al.* Comparison of urinary 6- β -cortisol and the erythromycin breath test as measures of hepatic

- P450IIIa (CYP3A) activity. *Clin Pharmacol Ther* 1992; 52: 265–273.
79. Hunt CM, Watkins PB, Saenger P *et al.* Heterogeneity of CYP3A isoforms metabolizing erythromycin and cortisol. *Clin Pharmacol Ther* 1992; 51: 18–23.
80. Schuetz EG, Schuetz JD, Grogan WM *et al.* Expression of cytochrome P450 3A in amphibian, rat and human kidney. *Arch Biochem Biophys* 1992; 294: 206–214.
81. Ohnhaus EE, Breckenridge AM, Park BK. Urinary excretion of 6 β -hydroxycortisol and the time course measurement of enzyme induction in man. *Eur J Clin Pharmacol* 1989; 36: 39–46.
82. Kovacs SJ, Martin DE, Everitt DE, Patterson SD, Jorkasky DK. Urinary excretion of 6 β -hydroxycortisol as an in vivo marker for CYP3A induction: Applications and recommendations. *Clin Pharmacol Ther* 1998; 63: 617–622.
83. Tran JQ, Kovacs SJ, McIntosh TS, Davis HM, Martin DE. Morning spot and 24-hour urinary 6 beta-hydroxycortisol to cortisol ratios: intraindividual variability and correlation under basal conditions and conditions of CYP 3A4 induction. *J Clin Pharmacol* 1999; 39: 487–494.
84. Mahmood I. Limited sampling model for the estimation of pharmacokinetic parameters in children. *Ther Drug Monit* 2000; 22: 532–536
85. Sun H, Pelsor F, Lazor J, Selen A, Lesko L. The reliability of population pharmacokinetic (PPK) approach in pediatric exclusivity claims (PEC). *Clin Pharmacol Ther* 2000; 67: 163 (Pill-86).
86. Burchell B, Coughtrie MW. Genetic and environmental factors associated with variation of human xenobiotic glucuronidation and sulfation. *Environ Health Perspect* 1997; 105 Suppl 4: 739–747.
87. Choonara IA, McKay P, Hain R, Rane A. Morphine metabolism in children. *Br J Clin Pharmacol* 1989; 28:599–604.
88. Anderson BJ, McKee AD, Holford NH. Size, myths and the clinical pharmacokinetics of analgesia in paediatric patients. *Clin Pharmacokinet* 1997; 33: 313–327.
89. Koren G, Butt W, Chinyanga H, Soldin S, Tan YK, Pape K. Postoperative morphine infusion in newborn infants: assessment of disposition characteristics and safety. *J Pediatrics* 1985; 107: 963–967.
90. McRorie TI, Lynn AM, Nespeca MK, Opheim KE, Slattery JT. The maturation of morphine clearance and metabolism. *Am J Dis Child* 1992; 146: 972–976.
91. Rollins DE, Bahr CV, Moldeus P, Rane A. Acetaminophen: Potentially toxic metabolite formed by human fetal and adult liver microsomes and isolated fetal liver cells. *Science* 1979; 205: 1414–1416.
92. Levy G, Khanna NN, Soda DM, Tsuzuki O, Stern L. Pharmacokinetics of acetaminophen in the human neonate: formation of acetaminophen glucuronide and sulfate in relation to plasma bilirubin concentration and D-glucuric acid excretion. *Pediatrics* 1975; 55: 818–825.
93. Kataria BK, Ved SA, Nicodemus HF *et al.* The pharmacokinetics of propofol in children using three different data analysis approaches. *Anesthesiology* 1994; 80: 104–122.
94. Ventatakrishnan K, von Moltke LL, Greenblatt DJ. Human drug metabolism and the cytochrome P450; Application and relevance on in vitro models; *J Clin Pharmacol* 2001; 41: 1149–1179.
95. Royal College of Paediatrics and Child Health: Ethics Advisory Committee. Guidelines for the ethical conduct of medical research involving children. *Arch Dis Child* 2000; 82: 177–182.
96. Lacroix D, Sonnier M, Moncion A, Cheron G, Cresteil T. Expression of CYP3A in the human liver. Evidence that the shift between CYP3A7 and CYP3A4 occurs immediately after birth. *Eur J Biochem* 1997; 247: 625–634.
97. Treluyer JM, Cheron G, Sonnier M, Cresteil T. Cytochrome P-450 expression in sudden infant death syndrome. *Biochem Pharmacol* 1996; 52: 497–504.
98. Johnson TN, Tanner MS, Taylor CT, Tucker GT. Enterocytic CYP3A4 in a paediatric population; developmental changes and the effects of celiac disease and cystic fibrosis. *Br J Clin Pharmacol* 2001; 51: 451–460.
99. Ueng YF, Kuwabara T, Chun YJ, Guengerich FP. Cooperativity in oxidations catalysed by cytochrome P450 3A4. *Biochemistry* 1997; 36: 370–381.
100. Kenworthy KE, Bloomers JC, Clarke KE, Houston JB; CYP3A4 drug interactions: correlation of 10 in vitro probe substrates. *Br J Clin Pharmacol* 1999; 48: 716–727.
101. Redfern M, Keeling J, Powell. The report of the Royal Liverpool Children's inquiry, HMSO 2000. Also available at www.rclinqury.org.uk/
102. Berthou F, Flinois JP, Ratanasavanh D, Beaune P, Riche C, Guillouzo A. Evidence for the involvement of several cytochromes P-450 in the first steps of caffeine metabolism by human liver microsomes. *Drug Metab Dispos* 1991; 19: 561–7.
103. Rane A, Phenotyping of drug metabolism in infants and children: Potentials and Problems. *Pediatrics* 1999; 104: 640–643.
104. Peng D, Birgersson C, von Bahr C, Rane A. Polymorphic acetylation of 7-amino-clonazepam in human liver cytosol. *Pediatr Pharmacol* 1984; 4: 155–159.
105. Murry GI, Barnes TS, Sewell HF, Ewan SWB, Melvin WT, Burke MD. The immunocytochemical localisation and distribution of cytochrome P-450 in normal human hepatic and extrahepatic tissues with a monoclonal antibody to human cytochrome P-450. *Br J Clin Pharmacol* 1988; 25: 465–475.
106. Strassburg CP, Strassburg A, Kneip S *et al.* Developmental aspects of human hepatic drug glucuronidation in young children and adults. *Gut* 2001; 50: 259–265.
107. Hines RN, McCarver DG. The ontogeny of human drug-metabolizing enzymes: Phase 1 oxidative enzymes. *J Pharmacol Exp Ther* 2002; 300: 355–360.
108. McCarver DG, Hines RN. The ontogeny of human drug-metabolizing enzymes: Phase 2 conjugation enzymes and regulatory mechanisms. *J Pharmacol Exp Ther* 2002; 300: 361–366.
109. Cresteil T, Beaune P, Kremers P, Celier C, Guengerich FP, Leroux JP. Immunoquantification of epoxide hydrolase and cytochrome P-450 isozymes in fetal and adult human liver microsomes. *Eur J Biochem* 1985; 151: 345 - 350.
110. Yang H-YL, Namkung MJ, Juchau MR. Expression of functional cytochrome P4501A1 in human embryonic hepatic tissue during organogenesis. *Biochem Pharmacol* 1995; 49: 717–726.
111. Hakkola J, Pasanen M, Purkunen R *et al.* Expression of xenobiotic-metabolizing cytochrome P450 forms in human adult and fetal liver. *Biochem Pharmacol* 1994; 48: 59–64.

112. Ratanasavanh D, Beaune P, Morel F, Flinois JP, Guengerich FP, Guillouzo A. Intralobular distribution and quantitation of cytochrome P450 enzymes in human liver as a function of age. *Hepatology* 1991; 13: 1142–1151.
113. Tateishi T, Nakura H, Asoh M *et al.* A comparison of hepatic cytochrome P450 protein expression between infancy and post infancy. *Life Sci* 1997; 61: 2567–2574.
114. Treluyer JM, Gueret G, Cheron G, Sonnier M, Cresteil T. Developmental expression of CYP2C and 2C dependent activities in the human liver: in vivo/in vitro correlation and inducibility. *Pharmacogenetics* 1997; 7: 441–452.
115. Treuyer JM, Benech H, Colin I, Pruvost A, Cheron G, Cresteil T. Ontogenesis of CYP2C-dependent arachidonic acid metabolism in the human liver: relationship with sudden infant death syndrome. *Pediatric Res* 2000; 47: 677–683.
116. Jacqz-Aigrain E, Cresteil T. Cytochrome P450-dependent metabolism of dextromethorphan: fetal and adult studies. *Dev Pharmacol Ther* 1992; 18: 161–8.
117. Treluyer JM, Jacqz-Aigrain E, Alvarez F, Cresteil T. Expression of CYP2D6 in developing human liver. *Eur J Biochem* 1991; 202: 583–588.
118. Jacqz-Aigrain E, Funck-Brentano C, Cresteil T. CYP2D6- and CYP3A-dependent metabolism of dextromethorphan in humans. *Pharmacogenetics* 1993; 3: 197–204.
119. Carpenter SP, Lasker JM, Raucy JL. Expression, induction and catalytic activity of the ethanol-inducible cytochrome P450 (CYP2E1) in human fetal liver and hepatocytes. *Mol Pharmacol* 1996; 49: 260–268.
120. Vieira I, Sonnier M, Cresteil T. Developmental expression of CYP2E1 in the human liver. Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 1996; 238: 476–483.
121. Boutelet-Bochan H, Huang Y, Juchau MR. Expression of CY2E1 during embryogenesis and fetogenesis in human cephalic tissues: Implications for the fetal alcohol syndrome. *Biochem Biophys Res Commun* 1997; 238: 443–447.
122. Gu J, Su T, Chen Y, Zhang Q-Y, Ding X. Expression of biotransformation enzymes in human fetal olfactory mucosa: Potential roles in developmental toxicity. *Toxicol Appl Pharmacol* 2000; 165: 158–162.
123. Treluyer JM, Rey E, Pons G, Cresteil T. Evidence of impaired cisapride metabolism in neonates. *Br J Clin Pharmacol* 2001; 52: 419–425.
124. Kawade N, Onishi S. The prenatal and postnatal development of UDP-glucuronyltransferase activity towards bilirubin and the effect of premature birth on this activity in the human liver. *Biochem J* 1981; 196: 257–260.
125. Leakey JE, Hume R, Burchell B. Development of multiple activities of UDP-glucuronyltransferase in human liver. *Biochem J* 1987; 243: 859–861.
126. Burchell B, Coughtrie M, Jackson M *et al.* Development of human liver UDP-glucuronosyltransferases. *Dev Pharmacol Ther* 1989; 13: 70–77.
127. Pacifici GM, Franchi M, Giuliani L, Rane A. Development of the glucuronyltransferase and sulphotransferase towards 2-naphthol in human fetus. *Dev Pharmacol Ther* 1989; 14: 108–114.
128. Pacifici GM, Sawe J, Kager L, Rane A. Morphine glucuronidation in human fetal and adult liver. *Eur J Clin Pharmacol* 1982; 22: 553–558.
129. Coughtrie MW, Burchell B, Leakey JE, Hume R. The inadequacy of perinatal glucuronidation: immunoblot analysis of the developmental expression of individual UDP-glucuronosyltransferase isoenzymes in rat and human liver microsomes. *Mol Pharmacol* 1988; 34: 729–735.
130. Wright MC, Edwards RJ, Pimenta M, Ribeiro V, Ratra GS. Developmental changes in the constitutive and inducible expression of cytochrome P450 3A2. *Biochem Pharmacol* 1997; 54: 841–846.
131. Johnson TN, Tanner MS, Tucker GG. Ontogeny of enterocytic and hepatic cytochrome P450 3A in the rat. *Biochem Pharmacol* 2000; 60: 1601–1610.
132. Rich KJ, Boobis AR. Expression and inducibility of P450 enzymes during liver ontogeny. *Micros Res Tech* 1997; 39: 424–435.
133. Mahnke A, Strotkamp D, Ross PH, Hanstein WG, Charbot GC, Neff P. Expression and inducibility of cytochrome P450 3A9 (CYP3A9) and other members of the CYP3A subfamily in rat liver. *Arch Biochem Biophys* 1997; 337: 62–68.
134. Shimada T, Guengerich FP. Participation of a rat liver cytochrome P450 induced by pregnenolone 16 alpha-carbonitrile and other compounds in the 4-hydroxylation of mephenytoin. *Mol Pharmacol* 1985; 28: 215–219.
135. Shimada T, Misono KS, Guengerich FP. Human liver microsomal cytochrome P-450 mephenytoin 4-hydroxylase, a prototype of genetic polymorphism in oxidative drug metabolism: purification and characterization of two similar forms involved in the reaction. *J Biol Chem* 1986; 261: 909–921.
136. Bonate PL, Howard D. Critique of prospective allometric scaling: does the emperor have clothes? *J Clin Pharmacol* 2000; 40: 335–340.
137. Cummings AG, Steele TW, Labrooy JT, Shearman DJC. Maturation of the rat small intestine at weaning: changes in epithelial cell kinetics, bacterial flora, and mucosal immune activity. *Gut* 1988; 29: 1672–1679.
138. Menard D. Development of human intestinal and gastric enzymes. *Acta Paediatrica*. 1994; 405: S1–6.
139. Zeugge J, Schneider G, Coassolo P, Lave T. Prediction of hepatic metabolic clearance. Comparison and assessment of prediction models. *Clin Pharmacokinet* 2001; 40: 553–563.
140. Chappell WR, Mordenti J. Extrapolation of toxicological and pharmacological data from animals to man. *Adv Drug Res* 1991; 20: 2–116.
141. Lave T, Dupin S, Scmitt C, Chou RC, Jaek D, Coassolo P. Integration of in vitro data into allometric scaling to predict hepatic metabolic clearance in man: application in 10 extensively metabolised drugs. *J Pharm Sci* 1997; 86: 584–590.
142. Ekins S, Ring B, Grace J, McRobie-Bell DJ, Wrighton SA. Present and future in vitro approaches for drug metabolism. *J Pharmacol Toxicol Meth* 2000; 44: 313–324.
143. Ekins S, Waller CL, Swaan PW, Cruciani G, Wrighton SA, Wikel JH. Progress in predicting human ADME parameters in silico. *J Pharmacol Toxicol Methods* 2000; 44: 251–272.
144. Tooley A, Rostami-Hodjegan A, Lennard MS, Tucker GT. Prediction of the interindividual variability; of in vivo metabolic interactions between ritonavir and methadone: acute versus long term effects. *Clin Pharmacol Ther* 2000; 68: 216.

145. Yang JS, Rostami-Hodjegan A, Tucker GT. Prediction of ketoconazole interaction with midazolam, alprazolam and triazolam: incorporating population variability. *Br J Clin Pharmacol* 2001; 48:4P.
146. Yang JS, Rostami-Hodjegan A, Tucker GT. Prediction of fluconazole interaction with midazolam and triazolam: incorporating population variability. *Br J Clin Pharmacol* 2001; 48:4P.
147. Yang JS, Rostami-Hodjegan A, Tucker GT. Prediction of ritonavir interaction with sildenafil (viagra): incorporating population variability. *Br J Clin Pharmacol* 2002 (in press).